

Employment of methyl 2-pyridyl ketone oxime in manganese non-carboxylate chemistry: $\text{Mn}^{\text{II}}_2\text{Mn}^{\text{IV}}$ and $\text{Mn}^{\text{II}}_2\text{Mn}^{\text{III}}_6$ complexes

Constantinos C. Stoumpos,^a Theocharis C. Stamatatos,^b Harikleia Sartzi,^a Olivier Roubeau,^c Anastasios J. Tasiopoulos,^d Vassilios Nastopoulos,^a Simon J. Teat,^e George Christou*^b and Spyros P. Perlepes*^a

^a *Department of Chemistry, University of Patras, 265 04 Patras, Greece. E-mail: perlepes@patreas.upatras.gr; Tel: +30 2610 997146.*

^b *Department of Chemistry, University of Florida, Gainesville, Florida, 32611-7200, USA. E-mail: christou@chem.ufl.edu; Tel: +1 352 392 8314.*

^c *CNRS, UPR 8641, Centre de Recherche Paul Pascal (CRPP), Equipe "Materiaux Moléculaires Magnétiques", 115 avenue du Dr. Albert Schweitzer, Pessac, F-33600, France.*

^d *Department of Chemistry, University of Cyprus, 1678 Nicosia, Cyprus.*

^e *Advanced Light Source, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Mail Stop 2-400, Berkeley, CA 94720, USA.*




Fig. S1 The out-of-phase (χ_M'') vs. T ac susceptibility signals for **3** in a 3.5 G field oscillating at the indicated frequencies.




Fig. S2 The out-of-phase (χ_M'') vs. T ac susceptibility signals for **4** in a 3.5 G field oscillating at the indicated frequencies.