

Electronic Supplementary Information for:

Red-light-emitting electrochemical cell using a polypyridyl iridium (III) polymer

José Lorenzo Rodríguez-Redondo,^a Rubén D. Costa,^b Enrique Ortí,^b Angela Sastre-Santos,^a Henk J. Bolink,^{*b} and Fernando Fernández-Lázaro^{*a}

^a *División de Química Orgánica, Instituto de Bioingeniería, Universidad Miguel Hernández, Elche 03202 (Alicante), Spain. E-mail: fdofdez@umh.es*

^b *Instituto de Ciencia Molecular, Universidad de Valencia, PO Box 22085, ES-46071, Valencia, Spain. E-mail: henk.bolink@uv.es*

Contents:

CHARACTERIZATION BY NMR SPECTROSCOPY

Figure S1 — Partial ^1H -NMR of a) monomer 7 and b) polymer 2 .-----	S3
Figure S2 — DQF-COSY of monomer 7 -----	S4
Figure S3 — HSQC of monomer 7 -----	S5
Figure S4 — HSQC of complex 1 -----	S6

CHARACTERIZATION BY NMR SPECTROSCOPY:




Figure S1. a) ¹H-NMR in CDCl₃ of the monomer **7** and b) ¹H-NMR in CD₂Cl₂ of the polymer **2**.




Figure S2. DQF-COSY of monomer 7.




Figure S3. HSQC of monomer 7.




Figure S4. HSQC of complex 1.