

Isomerization of the Hydride Complexes
 $[\text{HFe}_2(\text{SR})_2(\text{PR}_3)_x(\text{CO})_{6-x}]^+$ ($x = 2, 3, 4$) Relevant to the Active Site
Models for the [FeFe]-Hydrogenases

Bryan E. Barton, Giuseppe Zampella, Aaron K. Justice, Luca De Gioia*, Thomas B.
Rauchfuss*, Scott R. Wilson

Supporting Information




Figure S1. ^1H NMR (500 MHz) spectra of a CD_2Cl_2 solution of **1**, $\text{Fe}_2(\text{edt})(\text{CO})_4(\text{dppv})$, after protonation with 3 equiv $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ at -80°C , then warmed to recorded temperatures. Spectrum recorded at 20°C is after 48 h at room temperature.




Figure S2. ^1H NMR (500 MHz) spectra of a CD_2Cl_2 solution of **2**, $\text{Fe}_2(\text{pdt})(\text{CO})_4(\text{dppv})$, after protonation with 3 equiv $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ at $-80\text{ }^\circ\text{C}$, then warmed to recorded temperatures. Spectrum recorded at $20\text{ }^\circ\text{C}$ is after 16 h at room temperature.




Figure S3. Protonation of **2**, $\text{Fe}_2(\text{pdt})(\text{CO})_4(\text{dppv})$, with 3 equiv $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ at -80 °C, then warmed to recorded temperatures. $^{31}\text{P}\{^1\text{H}\}$ NMR (202 MHz) acquired with -50 °C ^1H NMR above shows the growth of signals at 97 and 72 ppm ($[\text{2H-Term}]^+$, for terminal hydride derivative wherein dppv is apical, basal) and a signal at 89 ppm ($[\text{2H-Term}^*]^+$, terminal hydride with hydride on dibasal-dppv side). When sample is warmed to -30 °C, new signals are observed at 92 and 95 ppm ($[\text{2H-A}]^+$ - is bridging hydride of apical,basal-dppv). When sample is warmed to room temperature, only one signal is observed at 86 ppm ($[\text{2H-B}]^+$ - thermodynamic product with dppv as dibasal (triplet in high-field ^1H NMR). At low temperature, signals at 96 and 82 ppm are due to remaining starting material.




Figure S4. ^1H NMR (500 MHz) spectrum of a CD_2Cl_2 solution of **4**, $\text{Fe}_2(\text{pdt})(\text{CO})_3(\text{PMe}_3)(\text{dppv})$, after protonation with $\text{HBF}_4 \cdot \text{Et}_2\text{O}$ at $-90\text{ }^\circ\text{C}$. Signals at -3, -4 ppm are assigned to isomers of the terminal hydride and appear as doublets with $J_{\text{PH}} \sim 75\text{ Hz}$. The intense signal at -14.3 ppm with $J_{\text{PH}} \sim 30\text{ Hz}$ is assigned to $[\mathbf{4H-A}]^+$. For spectra at higher temperature, see Figure S5.




Figure S5. ^1H NMR (500 MHz) spectra of CD_2Cl_2 solution of **4** $\text{Fe}_2(\text{pdt})(\text{CO})_3(\text{PMe}_3)(\text{dppv})$ after warming to various temperatures. Labeling is as follows (for peaks observed at $-30\text{ }^\circ\text{C}$): δ -14.4 (dt) is $[\mathbf{4H-A}]^+$, δ -14.7 (td) is $[\mathbf{4H-B}^*]^+$, δ -15.3 (td) is $[\mathbf{4H-B}]^+$ (for peaks observed at $23\text{ }^\circ\text{C}$): δ -14.3 (q) is $[\mathbf{4H-C}]^+$, δ -14.8 (td) is $[\mathbf{4H-D}]^+$. Peak at δ -16.4 is unknown.




Figure S6. ^1H NMR (500 MHz) spectra of CD_2Cl_2 solution of **3**, $\text{Fe}_2(\text{edt})(\text{CO})_3(\text{PMe}_3)(\text{dppv})$, after protonation with $[\text{H}(\text{Et}_2\text{O})_2]\text{BAr}^{\text{F}}_4$ at $-90\text{ }^\circ\text{C}$ and recorded at various temperatures. At $-75\text{ }^\circ\text{C}$, a doublet is the sole species present in the ^1H NMR spectrum, $J_{\text{PH}} \sim 30\text{ Hz}$ (indicating a single P atom *cis* to the hydride). Upon warming to $-30\text{ }^\circ\text{C}$, two new triplets are observed in the ^1H NMR, $J_{\text{PH}} \sim 25\text{ Hz}$ (indicating two P atoms *cis* to the hydride). Upon equilibration of $[\text{3H}]^+$ at room temperature for several months, the ^1H NMR displays a triplet and a triplet of doublets, and possibly one more signal with unresolved coupling under the major resonance at δ -16.8. As seen for $[\text{4H}]^+$, the unstable rotamer $[\text{3H-B}^*]^+$ is observed at δ -16.8.




Figure S7. $^{31}\text{P}\{^1\text{H}\}$ NMR spectra of CD_2Cl_2 solution of **3**, $\text{Fe}_2(\text{edt})(\text{CO})_3(\text{PMe}_3)(\text{dppv})$, after protonation with $[\text{H}(\text{Et}_2\text{O})_2]\text{BAr}^{\text{F}}_4$ at $-90\text{ }^\circ\text{C}$ and recorded at various temperatures. At $-75\text{ }^\circ\text{C}$, the $^{31}\text{P}\{^1\text{H}\}$ spectrum shows mostly one isomer, with two signals in the dppv region (apical-basal) and one signal in the PMe_3 region. At $-30\text{ }^\circ\text{C}$, the same peaks remain, smaller broad peaks are due to slight remainder of starting material. Upon equilibration of $[\mathbf{3H}]^+$ at room temperature for several months, the $^{31}\text{P}\{^1\text{H}\}$ NMR spectrum displays in the dppv region a set of equally intense peaks at 92, 93 ppm (apical-basal, major isomer), a single resonance at 89 ppm (dibasal, minor isomer), and in the PMe_3 region three peaks. The missing dppv signal for the third isomer wherein the dppv is possibly dibasal is not observed.

QuickTime™ and a
decompressor
are needed to see this picture.

Figure S8. High-field ^1H NMR spectra of a d^7 -DMF solution of $[\mathbf{4H}]\text{PF}_6$, **a)** recorded at 20 °C on 500 MHz instrument and **b)** after heating to 80 °C. The spectrum recorded at 120 °C (600 MHz) was after three days at room temperature, decomposition species are labeled: $\text{Fe}_2(\text{pdt})(\text{CO})_2(\text{dppv})_2$ (—) and unknown poly-phosphine species (—). At 80°C the peaks at –13.8 ppm [**4H-C**] and –14.5 [**4H-D**] are broadened. At 120 °C, [**4H-B**] and [**4H-C**] are labeled with (—).




Figure S9. $^{31}\text{P}\{^1\text{H}\}$ NMR (600 MHz, CD_2Cl_2) spectrum of $[\text{HFe}_2(\text{edt})(\text{CO})_2(\text{dppv})_2]^+$ recorded at -35°C . Peaks corresponding to $[\mathbf{5H}\text{-Term}]^+$ are at 99, 91, 84, 74 ppm. Remaining signals are due to $[\mathbf{5H}\text{-A}]^+$ (97, 87 ppm) and $\mathbf{5}$ (92, 93 ppm).




Figure S10. Difference in energy for isomers for $[3H]^+$ and $[4H]^+$, $[HFe_2(xdt)(CO)_3(PMe_3)(dppv)]BF_4$, as calculated experimentally.




Figure S11. Plot monitoring the concentration of $[4\text{H-B}]^+$, $[4\text{H-C}]^+$ and $[4\text{H-D}]^+$ as a function of time. Data was collected using ^1H NMR spectroscopy.




Table S1. Crystallographic data for $[3H-B]^+$, $[HFe_2(edt)(CO)_3(dppv)(PMe_3)]PF_6$, including selected lengths (Å) and angles (°).

Fe(1) – Fe(2)	2.5744 (5)	Fe(1) – S(1) – Fe(2)	69.355 (17)
Fe(1) – S(1)	2.2581 (7)	Fe(1) – Fe(2) – S(1)	55.164 (17)
Fe(1) – S(2)	2.2794 (6)	Fe(2) – Fe(1) – P(1)	148.911 (18)
Fe(2) – S(1)	2.2668 (7)	Fe(2) – Fe(1) – P(2)	113.20 (2)
Fe(2) – S(2)	2.2760 (6)	Fe(2) – Fe(1) – C(1)	111.37 (6)
Fe(1) – P(1)	2.2137 (6)	P(1) – Fe(1) – P(2)	86.80 (2)
Fe(1) – P(2)	2.2282 (6)	P(1) – Fe(1) – C(1)	90.78 (6)
Fe(1) – C(1)	1.765 (2)	P(2) – Fe(1) – C(1)	91.10 (6)
Fe(2) – P(3)	2.2549 (7)	Fe(1) – Fe(2) – C(2)	112.44 (7)
Fe(2) – C(2)	1.790 (2)	Fe(1) – Fe(2) – C(3)	141.78 (7)
Fe(2) – C(3)	1.771 (2)	Fe(1) – Fe(2) –	109.54 (2)

		P(3)	
Fe(1) – H(1)	1.69 (2)	C(3) – Fe(2) – C(2)	95.45 (10)
Fe(2) – H(1)	1.64 (2)	C(3) – Fe(2) – P(3)	94.15 (7)
C(1) – O(1)	1.142 (2)	C(2) – Fe(2) – P(3)	92.44 (7)
C(2) – O(2)	1.140 (3)	Fe(1) – H(1) – Fe(2)	99.99
C(3) – O(3)	1.148 (3)		

Complex	[Fe ₂ (edt)(μ-H)(CO) ₃ (PMe ₃)(dppv)]PF ₆
Chemical formula	C ₃₄ H ₃₆ F ₆ Fe ₂ O ₃ P ₄ S ₂
Temperature (K)	193 (2)
Crystal size (mm ³)	0.40 x 0.24 x 0.16
Crystal system	Triclinic
Space group	P-1
a (Å)	10.818 (2)
b (Å)	11.253 (2)
c (Å)	16.208 (3)
α (°)	78.473 (3)
β (°)	86.415 (3)
γ (°)	81.525 (3)
V (Å ³)	1911.1 (6)
Z	2
Density calcd (Mg m ⁻³)	1.575
μ (Mo Kα, mm ⁻¹)	0.71073
max./min. trans'n	0.8572 / 0.6869
reflections meas'd/Indep.	23821 / 9184
data/restraints/parameters	9184 / 0 / 467
GOF on F ²	1.024
Rint	0.0286
R1 [I > 2σ] (all data) ^a	0.0322 (0.0485)
wR2 [I > 2σ] (all data) ^b	0.0786 (0.0850)
max. peak/hole (e ⁻ /Å ³)	0.392 / -0.374

