

Supplementary Information

Luminescence of Five-coordinated Nickel(II) Complexes with Substituted- 8-Hydroxyquinolines and Macroyclic Ligands

M. Dolores Santana,^a Rocío García-Bueno,^a and Gabriel García,^a José Pérez,^b Luis García,^b Miguel Monge,^c and Antonio Laguna^d*

^a Departamento de Química Inorgánica, Universidad de Murcia, 30071 Murcia, Spain.

^b Departamento de Ingeniería Minera, Geológica y Cartográfica. Área de Química Inorgánica, Universidad Politécnica de Cartagena, E-30203 Cartagena, Spain.

^c Departamento de Química, Complejo Científico-Técnico, Universidad de La Rioja, 26004 Logroño, Spain.

^d Departamento de Química Inorgánica, ICMA, Universidad de Zaragoza-CSIC, 50009-Zaragoza, Spain.




Figure 1S. Magnitude $\{^1\text{H}-^1\text{H}\}$ COSY spectrum of complex **4b** at 200MHz at 21 °C in $(\text{CD}_3)_2\text{CO}$ solution. Only the region relevant to the assignment of quinolinolate resonances is shown in the top trace.




Figure 2S. Magnitude $\{^1\text{H}-^1\text{H}\}$ COSY spectrum of complex **7a** at 200MHz at 21 °C in $(\text{CD}_3)_2\text{CO}$ solution. Only the region relevant to the assignment of quinolinolate resonances is shown in the top trace.