

Two unprecedented NLO-active coordination polymers constructed by a semi-rigid tetrahedral linker

Li-Li Liang, Shi-Bin Ren, Jun Zhang, Yi-Zhi Li, Hong-Bin Du* and Xiao-Zeng You
*State Key Laboratory of Coordination Chemistry, School of Chemistry and Chemical
Engineering, Nanjing University, Nanjing 210093, China. E-mail: [hbdu@nju.edu.cn](mailto:hbd@nju.edu.cn)*

Supporting Information

Contents:

1. Figure S1. The asymmetric unit of complex **1**.
2. Figure S2. Coordination environments of binuclear Zn₂ units in **1**.
3. Figure S4. PXRD patterns for **1**.
4. Figure S5. PXRD patterns for **2**.




Figure S1. Perspective view of the asymmetric units of complex **1**.




Figure S2. The different coordination environments of Zn1Zn2 and Zn3Zn4 binuclear Zn₂ units in **1**.




Figure S3. PXRD patterns for **1**: (a) simulated (black), (b) as-synthesized (red), (c) heated at 100°C for 2h (magenta), (d) heated at 200°C for 2h (blue), (e) heated at 300°C for 2h (green) in air.




Figure S4. PXRD patterns for **2**: (a) simulated (black), (b) as-synthesized (red).