

Supporting Information

Two additive-induced isomeric aluminoborates templated by methylamine

Gao-Juan Cao,^a Jian Lin,^c Jin-Yun Wang,^a Shou-Tian Zheng,^a Wei-Hui Fang^a and Guo-Yu Yang^{a,b}

^aState Key Laboratory of Structural Chemistry, Fujian Institute of Research on the Structure of Matter, Chinese Academy of Sciences, Fuzhou, 350002, P.R. China. E-mail: ygy@fjirsm.ac.cn; Fax: (+86) 591-8371-0051

^bDepartment of Chemistry, Beijing Institute of Technology, Beijing 100081, P. R. China.

^cXiamen University, Xiamen 361005, P. R. China.

Fig. S1 IR spectra of **1** and **2**, respectively.

Fig. S2 The asymmetric unit of compound **1**.

Fig. S3 View of the coordination environments of all the B and Al atoms in compound **1**.

Fig. S4 View of the anionic framework of **1**, showing cone-shaped 10-ring channels along the [110] (**a**) and [-110] (**b**) directions.

Fig. S5 View of the unclosed 10-ring channels along the [110] (**a**) and [-110] (**b**) directions in compound **1**.

Fig. S6 View of the templates and guest water molecules located in the 10-ring channel, interacted with the walls via hydrogen bonds in compound **1**.

Fig. S7 IR spectra (**a**) and powder X-ray diffraction patterns (**b**) of compound **1** (black line), Na⁺- (red line), K⁺- (blue line) and Ag⁺-exchange (cyan) samples.

Fig. S8 View of the coordination environments of all the B and Al atoms in compound **2**.

Fig. S9 IR spectra (**a**) and powder X-ray diffraction patterns (**b**) of compound **2** (black line), Na⁺- (red line), K⁺- (blue line) and Ag⁺-exchange (cyan) samples.

Fig. S10 View of three different shaped 11-ring channels in **2**, propagated along the [001], [110] and [112] directions, respectively.

Fig. S11 View of the 10-ring channels in **2**, running along the [101] direction.




Fig. S1 IR spectra of **1** and **2**, respectively.




Fig. S2 The asymmetric unit of compound **1**.




Fig. S3 View of the coordination environments of all the B and Al atoms in compound **1**. Symmetry codes for **1**: #A x, -y+0.5, -z+0.25, #B x-1, -y+1, -z, #C -x+2, -y, z, #D -x+1, y+1, -z, #E -x+2, y+1, -z, #F -x+3, -y+1, z, #G x-1, -y+2, -z.




Fig. S4 View of the anionic framework of **1**, showing cone-shaped 10-ring channels along the [110] (**a**) and [-110] (**b**) directions in compound **1**. Color code: B₂O₄, yellow; B₁O₃, B₃O₃, purple; AlO₄, green.




Fig. S5 View of the unclosed 10-ring channels along the [110] (a) and [-110] (b) directions in compound 1. Color code: B2, yellow; B1, B3, purple; Al, green.




Fig. S6 View of the templates and guest water molecules located in the 10-ring channel, interacted with the walls via hydrogen bonds in compound 1.




Fig. S7 IR spectra (a) and powder X-ray diffraction patterns (b) of compound 1 (black line), Na^+ - (red line), K^+ - (blue line) and Ag^+ -exchange (cyan) samples.




Fig. S8 View of the coordination environments of all the B and Al atoms in compound 2. Symmetry codes for 2: #A -x+1, y, -z+3, #B -x, y, -z+2, #C -x+1/2, y+1/2, -z+2, #D x-1/2, y+1/2, z.




Fig. S9 IR spectra (a) and powder X-ray diffraction patterns (b) of compound 2 (black line), Na⁺ (red line), K⁺ (blue line) and Ag⁺-exchange (cyan) samples.




Fig. S10 View of three different shaped 11-ring channels in **2**, propagated along [001], [110] and [112] directions. Colour code: B_3O_4 , yellow; B_1O_3 , B_2O_3 , purple; AlO_4 , green.




Fig. S11 View of the 10-ring channels in **2**, running along [101] direction. Colour code: B_3O_4 , yellow; B_1O_3 , B_2O_3 , purple; AlO_4 , green.