

Supplementary Information

**Ferrocenyl catechols: Synthesis, oxidation chemistry and anti-proliferative effects on
MDA-MB-231 breast cancer cells**

Yong Leng Kelvin Tan,^{a,b,c} Pascal Pigeon,^{a,b} Siden Top,^{a,b} Eric Labbé,^d Olivier Buriez,^d Elizabeth A. Hillard,*^{a,b} Anne Vessières,^{a,b} Christian Amatore,^d Weng Kee Leong^e and Gérard Jaouen*^{a,b}

^a ENSCP Chimie ParisTech, Laboratoire Charles Friedel (LCF), 75005 Paris, France

^b CNRS, UMR 7223, 75005 Paris, France

^c Hwa Chong Institution, 661 Bukit Timah Road, Singapore 269734, Singapore

^d Ecole Normale Supérieure, Département de Chimie, 24 rue Lhomond, 75231 Paris cedex 05, France

^e Division of Chemistry and Biological Chemistry, School of Physical and Mathematical Sciences, Nanyang Technological University, 21 Nanyang Link, Singapore 637371, Singapore

Figure S1. Cyclic voltammogram for **1a**, **2c**, **3c** and **3d**.

Figure S2. ¹H NMR spectrum for **8a**.

Figure S3. ¹H NMR spectrum for **8b**.

Figure S4. ¹H NMR spectrum for **8c**.

Figure S5. ¹H NMR spectrum for **8d**.




Figure S1. Cyclic voltammogram for **2c** (1 mM in 0.1 M $\text{Bu}_4\text{NPF}_6/\text{CH}_2\text{Cl}_2$) in the absence (solid line) and presence (dashed line) of lutidine in 1:50 volume ratio. Scan rate 0.1 V s^{-1} . Pt electrode of 0.5 mm diameter.




Figure S2. ¹H NMR spectrum for **8a**.




Figure S3. ¹H NMR spectrum for **8b**.




Figure S4. ^1H NMR spectrum for **8c**.




Figure S5. ^1H NMR spectrum for **8d**.