

Supplementary Information

Allyl Strontium Compounds: Synthesis, Molecular Structure and Properties

Phillip Jochmann,^a Julien P. Davin,^a Stefanie Maslek,^a Thomas P. Spaniol,^a Yann Sarazin,^b Jean-Francois Carpentier*^b and Jun Okuda*^a

^a Institute of Inorganic Chemistry, RWTH Aachen University, Landoltweg 1, D-52056 Aachen, Germany. Fax: +49 241 80 92 644; Tel: +49 241 80 94 645; E-mail: jun.okuda@ac.rwth-aachen.de

^b Laboratoire Catalyse et Organométalliques, CNRS - Université de Rennes 1 - Sciences Chimiques de Rennes (UMR 6226), Campus de Beaulieu, 35042, Rennes Cedex, France. Fax:(+33 2 2323 6939; Tel: +33 2 2323 5880; E-mail: sophie.guillaume@univ-rennes1.fr

Contents

NMR spectra of

[Sr(C ₃ H ₅) ₂] (1 ; <i>in situ</i>)	S2
K[Sr(C ₃ H ₅) ₃] (2)	S2




Figure S1. ^1H NMR spectrum of $[\text{Sr}(\text{C}_3\text{H}_5)_2]$ (**1**; *in situ*) in $d_8\text{-thf}$ (*) at 24 °C. § denotes resonances of poly(thf).




Figure S2. ^1H NMR spectrum of $[\text{K}^+ \{ \text{Sr}(\text{C}_3\text{H}_5)_2 \}]$ (**2**) in $d_8\text{-thf}$ (*) at 23 °C.

SM 10 VT.007.esp




Figure S3. ^1H NMR spectrum of $[\text{K}\{\text{Sr}(\text{C}_3\text{H}_5)_2\}]$ (**2**) in $\text{d}_8\text{-thf}$ (*) at -95°C .




Figure S4. ^{13}C NMR spectrum of $[\text{K}\{\text{Sr}(\text{C}_3\text{H}_5)_2\}]$ (**2**) in $\text{d}_8\text{-thf}$ (*) at 23°C .