Supporting Information

Mixed ligand palladium(II) complexes of 6-methoxy-2-oxo-1,2dihydroquinoline-3-carbaldehyde 4N-substituted thiosemicarbazones with triphenylphosphine co-ligand: synthesis, crystal structure and biological properties

Eswaran Ramachandran,^a Duraisamy Senthil Raja,^a Nattamai S. P. Bhuvanesh^b and Karuppannan Natarajan^{*a}

^aDepartment of Chemistry, Bharathiar University, Coimbatore, 641 046, India. ^bDepartment of Chemistry, Texas A&M University, College Station, TX77843, USA.

*Corresponding author. Tel.: +91 422 2428319; fax: +91 422 2422387.

E-mail address: *k_natraj6@yahoo.com*



Figure S1 Synchronous spectra of BSA (1 μ M) in the presence of increasing amounts of the complexes 1(A), 2 (B), 3 (C), 4 (D) and 5 (E) (0-25 μ M) in the wavelength difference of $\Delta\lambda = 15$ nm. Arrow shows the emission intensity changes upon the increasing concentration of complexes.



Figure S2 Synchronous spectra of BSA (1 μ M) in the presence of increasing amounts of the complexes 1(A), 2 (B), 3 (C), 4 (D) and 5 (E) (0-25 μ M) in the wavelength difference of $\Delta\lambda = 60$ nm. Arrow shows the emission intensity changes upon the increasing concentration of complexes.