

Supporting Information

Bifunctional 3D Cu-MOFs containing glutarates and bipyridyl ligands: selective CO₂ sorption and catalytic activity

**In Hong Hwang,^a Jeong Mi Bae,^a Wan-Seok Kim,^b Young Dan Jo,^a Cheal Kim,^{*a}
Youngmee Kim,^{*c} Sung-Jin Kim,^c and Seong Huh^{*b}**

^a*Department of Fine Chemistry, Seoul National University of Science & Technology, Seoul 139-743, Korea. E-mail: chealkim@seoultech.ac.kr; Fax: 82-2-973-9149; Tel: 82-2-970-6693*

^b*Department of Chemistry and Protein Research Center for Bio-Industry, Hankuk University of Foreign Studies, Yongin 449-791, Korea. E-mail: shuh@hufs.ac.kr; Fax: 82-31-330-4566; Tel: 82-31-330-4522*

^c*Department of Chemistry and Nano Science, Ewha Womans University, Seoul 120-750, Korea. E-mail: ymeekim@ewha.ac.kr; Fax: 82-2-3277-2384; Tel: 82-2-3277-3589*

Table S1. Recycle experiments for the MOF 1.

Entry	Period (d) ^a	Conversion (%)
1	2	100
2	2	100
3	2	100
4	2	100
5	2	100
6	2	100
7	2	100
8	2	100
9	2	100
10	2	100
11	2	100
12	2	100
13	2	100
14	2	100
15	2	100
16	2	100
17	2	100
18	2	100
19	2	100
20	2	100

^aThe conversion was monitored by GC/Mass and the internal standard was dodecane.