

Supporting Information

Ruthenium(II) Polypyridyl Complexes as Dual Inhibitors of Telomerase and Topoisomerase

Guoliang Liao,[‡] Xiang Chen,[‡] Jingheng Wu, Chen Qian, Yi Wang, Liang-Nian Ji and Hui Chao*

MOE Key Laboratory of Bioinorganic and Synthetic Chemistry, State Key Laboratory of Optoelectronic Materials and Technologies, School of Chemistry and Chemical Engineering, Sun Yat-Sen University, Guangzhou 510275, P. R. China.

E-mail: ceschh@mail.sysu.edu.cn

[‡]These authors contributed equally to this work.

Fig. S1 ES-MS spectrum of complex 1	S2
Fig. S2 ¹ H NMR spectrum of complex 1	S2
Fig. S3 ES-MS spectrum of complex 1 co-incubated with L-cysteine.....	S3
Fig. S4 ES-MS spectrum of complex 1 co-incubated with glutathione	S3
Fig. S5 ES-MS spectrum of complex 1 co-incubated with dithiothreitol.....	S4
Fig. S6 ES-MS spectrum of complex 2 co-incubated with L-cysteine.....	S4
Fig. S7 ES-MS spectrum of complex 2 co-incubated with glutathione	S5
Fig. S8 ES-MS spectrum of complex 2 co-incubated with dithiothreitol.....	S5
Fig. S9 ES-MS spectrum of complex 3 co-incubated with L-cysteine.....	S6
Fig. S10 ES-MS spectrum of complex 3 co-incubated with glutathione	S6
Fig. S11 ES-MS spectrum of complex 3 co-incubated with dithiothreitol.....	S7




Fig. S1 ES-MS spectrum of complex 1




Fig. S2 ^1H NMR spectrum of complex 1




Fig. S3 ES-MS spectrum of complex **1** after co-incubated with L-cysteine under 37 °C for 1 h then R. T. for 24 h.




Fig. S4 ES-MS spectrum of complex **1** after co-incubated with glutathione under 37 °C for 1 h then R. T. for 24 h.




Fig. S5 ES-MS spectrum of complex **1** after co-incubated with dithiothreitol under 37 °C for 1 h then R. T. for 24 h.




Fig. S6 ES-MS spectrum of complex **2** after co-incubated with L-cysteine under 37 °C for 1 h then R. T. for 24 h.




Fig. S7 ES-MS spectrum of complex **2** after co-incubated with glutathione under 37 °C for 1 h then R. T. for 24 h.




Fig. S8 ES-MS spectrum of complex **2** after co-incubated with dithiothreitol under 37 °C for 1 h then R. T. for 24 h.




Fig. S9 ES-MS spectrum of complex **3** after co-incubated with L-cysteine under 37 °C for 1 h then R. T. for 24 h.




Fig. S10 ES-MS spectrum of complex **3** after co-incubated with glutathione under 37 °C for 1 h then R. T. for 24 h.




Fig. S1 ES-MS spectrum of complex **3** after co-incubated with dithiothreitol under 37 °C for 1 h then R. T. for 24 h.