Supporting Information Exploration of the active center structure of nitrogen-doped graphene-based catalysts for oxygen reduction reaction

Linfei Lai,^{*a,b,c*} Jeffrey R. Potts,^{*b*} Da Zhan,^{*a*} Liang Wang,^{*a*} Chee Kok Poh,^{*c*} Chunhua Tang,^{*d*} Hao

Gong,^d Zexiang Shen,^a Jianyi Lin,^c Rodney S. Ruoff^{*b}

^a Division of Physics and Applied Physics, School of Physical and Mathematical Sciences, Nanyang Technological University, 637371, Singapore

^b Department of Mechanical Engineering and the Materials Science and Engineering Program, University of Texas at Austin, One University Station C2200, Austin, TX 78712, USA

^c Institute of Chemical and Engineering Sciences, A*STAR, 1 Pesek Road, Jurong Island, 627833, Singapore

^d Department of Materials Science and Engineering, National University of Singapore, 117576, Singapore



Figure S1. B1s spectra of BN-RG-O.



Figure S2. Linear sweep voltammograms and Koutecky-Levich plots for BN-RG-O with rotation rate from 100 to 3000 rpm in 0.1 M KOH.



Figure S3. Linear sweep voltammograms and Koutecky-Levich plots for N-RG-O 550°C with rotation rate from 100 to 3000 rpm in 0.1 M KOH.



Figure S4. Linear sweep voltammograms and Koutecky-Levich plots for N-RG-O 850°C with rotation rate from 100 to 3000 rpm in 0.1 M KOH.



Figure S5. Koutecky-Levich plots for N-RG-O 1000°C with rotation rate from 100 to 3000 rpm in 0.1 M KOH.



Figure S6. Linear sweep voltammograms and Koutecky-Levich plots for PANi-RG-O with rotation rate from 100 to 3000 rpm in 0.1 M KOH.



Figure S7. Linear sweep voltammograms and Koutecky-Levich plots for PANi-RG-O with rotation rate from 100 to 3000 rpm in 0.1 M KOH.



Figure S8. Linear sweep voltammograms of N-RG-O and BN-RG-O in 0.1 M KOH at a rotating rate of 2500 rpm.