

Supplementary information for:

Perfluorinated compounds in maternal serum and cord blood from selected areas of South Africa: results of a pilot study

Linda Hanssen,^{1,2} Halina Röllin,^{3,4} Jon Øyvind Odland,¹ Morten K. Moe,² Torkjel M Sandanger^{1,2}

¹Department of Community Medicine, University of Tromsø, 9037 Tromsø, Norway

²Norwegian Institute for Air Research, the Polar Environmental Centre, 9296 Tromsø, Norway

³Medical Research Council, Johannesburg, South Africa

⁴School of Health Systems and Public Health, University of Pretoria, Pretoria, South Africa

Table S1 Socioeconomic and housing characteristics of participants by site

Statistics	Urban N=16	Industrial N=12	Atlantic Ocean N=12	Mining N=12	Indian Ocean malaria N=12	Inland malaria N=12
Population group %						
B=African Black	B 94	B 92	B 27			
C=Coloured				B 100	B 100	B 100
W=White	C 6	W 8	C 73			
Marital status (%)						
M=married	M 29	M 9	M 20	M 20		
S=single	S 17	S 75	S 40	S 40	S 100	S 100
L/T=living together			L/T 30			
D=divorced	L/T 54	L/T 16	D 10	L/T 40		
Mean monthly income Rand (1US\$ = 7.2 R)	4166 (4440)	No data	3286 (2276)	2250(1838)	No data	968 (547)
Residence ownership %	19	92	80	60	100	100
Type of housing %						
H=house	H41	H67		H82	H50	H88
F=Flat	F53	S33	H100	S18	S50	S12
S=Substandard	S6					
Electrified %	100	83	91	40	9	0
Potable drinking water %	100	100	100	100	45	42




Figure S1. Chromatogram of unspecified branched (RT7.13) and linear isomers (RT 7.37) of PFOS.