

**Supporting information (SI) of the article entitled:**

**Dietary gallic acid and anthocyanins cytotoxicity on human fibrosarcoma HT1080 cells. A study on the mode of action.**

Kamila Filipiak<sup>a,b,c</sup>, Maria Hidalgo<sup>c</sup>, Jose Manuel Silvan<sup>c</sup>, Benjamin Fabre<sup>a</sup>, Rodrigo J Carbajo<sup>d</sup>, Antonio Pineda-Lucena<sup>d</sup>, Ana Ramos<sup>a</sup>, Beatriz de Pascual-Teresa<sup>a</sup>, Sonia de Pascual-Teresa<sup>c</sup>

<sup>a</sup>Departamento de Química y Bioquímica, Facultad de Farmacia, Universidad CEU San Pablo, Urbanización Monteprincipe, 28668 Madrid, Spain

<sup>b</sup>Department of Molecular Biology, Faculty of Biotechnology and Environment Sciences, The John Paul II Catholic University of Lublin, 20-718, Lublin, Poland

<sup>c</sup>Department of Metabolism and Nutrition, Institute of Food Science, Food Technology and Nutrition (ICTAN), Spanish National Research Council (CSIC), José Antonio Novais 10, 28040-Madrid, Spain

<sup>d</sup>Structural Biochemistry Laboratory, Advanced Therapies Department, Centro de Investigación Príncipe Felipe, C/ Eduardo Primo Yúfera 3, 46012 Valencia, Spain

**\*Correspondence:** Sonia de Pascual-Teresa, Department of Metabolism and Nutrition, ICTAN-CSIC, Jose Antonio Novais 10, Ciudad Universitaria, Madrid 28040, Spain.

Fax: +34915493627, e-mail: [s.depascualteresa@csic.es](mailto:s.depascualteresa@csic.es)

## SUPPORTING INFORMATION

### Table of Content

Computational material.....	S2
-----------------------------	----

#### Computational material

**Figure S1.** Most energetically favored binding mode of gallic acid in MMP-2.

