

LIPASE-MEDIATED EPOXIDATION UTILIZING UREA-HYDROGEN PEROXIDE IN ETHYL ACETATE

Emmanuel Ankudey, Horacio F. Olivo, and Tonya L. Peebles
Chemical and Biochemical Engineering and
Division of Medicinal and Natural Products Chemistry
The University of Iowa, Iowa City, Iowa 52242*

Electronic Supplementary Information

Cyclohexene Oxide (1)	
¹ H-NMR spectra	S-3
¹³ C-NMR spectra	S-4
Methylcyclohexene oxide (2)	
¹ H-NMR spectra	S-5
¹³ C-NMR spectra	S-6
Phenylcyclohexene oxide (3)	
¹ H-NMR spectra	S-7
¹³ C-NMR spectra	S-8
Cyclooctene oxide (4)	
¹ H-NMR spectra	S-9
¹³ C-NMR spectra	S-10
exo-Norbornene oxide (5)	
¹ H-NMR spectra	S-11
¹³ C-NMR spectra	S-12
alpha-Pinene oxide (6)	
¹ H-NMR spectra	S-13
¹³ C-NMR spectra	S-14
Hexene oxide (7)	
¹ H-NMR spectra	S-15
¹³ C-NMR spectra	S-16

Octene oxide (8)	
¹ H-NMR spectra	S-17
¹³ C-NMR spectra	S-18
Styrene oxide (9)	
¹ H-NMR spectra	S-19
¹³ C-NMR spectra	S-20
<i>alpha</i>-Methylstyrene oxide (10)	
¹ H-NMR spectra	S-21
¹³ C-NMR spectra	S-22
<i>beta</i>-Methylstyrene oxide (11)	
¹ H-NMR spectra	S-23
¹³ C-NMR spectra	S-24
Indene oxide (12)	
¹ H-NMR spectra	S-25
¹³ C-NMR spectra	S-26















































