

Supporting Information for:

Synthesis of γ -Lactols, γ -Lactones and 1,4-Monoprotected Succinaldehydes Under Moderately Concentrated Sunlight

Daniele Dondi^{a*}, Stefano Protti^b, Angelo Albini^b, Sonia Mañas Carpio^{c,d} and Maurizio Fagnoni^b

^aDepartment of General Chemistry, University of Pavia, V.le Taramelli 10, 27100 Pavia (Italy)

^bDepartment of Organic Chemistry, University of Pavia, V.le Taramelli 12, 27100 Pavia (Italy)

^cÁrea de Química Inorgánica, Facultad de Ciencias, Universidad de Almería, Almería, 04120, (Spain)

^d Organometallic and Photochemistry laboratory for sustainable Chemistry, Ciesol, Almería, 04120, (Spain)

*E-mail: dondi@unipv.it; Fax: +39382528544; Tel: +39382987344

Figure S1: UV spectra of BPSS 10^{-4} M (solid line) and 10^{-2} M (dotted line) in water.





Figure S2: Solar radiation during the three days when the sun-driven photocatalyzed reaction between isopropanol and maleic acid was carried out (mid October to mid November period, Table 1, entry 1).






Figure S3: Solar irradiation during an average day in Almeria (from end June to mid July) when the reactions reported in entries 3-7 were carried out.

