Electronic Supplementary Information (ESI)

Nanostructured molybdenum carbides supported on carbon

nanotubes as efficient catalysts for one-step hydrodeoxygenation and

isomerization of vegetable oils

Junxing Han,^a Jinzhao Duan,^a Ping Chen,^a Hui Lou*^a, Xiaoming Zheng^a and Haiping Hong^b

^a Key Lab of Applied Chemistry of Zhejiang Province, Department of Chemistry, Zhejiang University, Hangzhou, 310028, China. E-mail: hx215@zju.edu.cn; Fax: +86-571-88273283; Tel: +86-571-88273283.

^b Department of Material and Metallurgical, South Dakota School of Mines and Technology, Rapid City, 57701, USA.

Table of contents

Bright-field TEM image of prepared Mo ₂ C/CNTs catalyst	.Fig.	S1
SEM image of prepared 10% Mo ₂ C/CNTs catalyst	Fig.	S2
SEM image of prepared 20% Mo ₂ C/CNTs catalyst	Fig.	S 3
SEM image of prepared 40% Mo ₂ C/CNTs catalyst	Fig.	S4
Distribution of fatty acids in vegetable oils	Table	e S 1



Fig. S1 Bright-field TEM image of prepared Mo₂C/CNTs catalyst.



Fig. S2 SEM image of prepared 10%Mo₂C/CNTs catalyst.



Fig. S3 SEM image of prepared 20%Mo₂C/CNTs catalyst.



Fig. S4 SEM image of prepared 40% Mo₂C/CNTs catalyst.

Fatty Acids	Linoleic	Linolenic	Oleic	Palmitic	Stearic	others			
Maize Oil	55.9	1.2	28.4	10.8	1.8	1.9			
Olive Oil	5.6	0.7	77.5	10.5	3.2	2.5			
Rapeseed Oil	21.3	7.4	59.5	6.6	2.3	2.9			
Soya Bean Oil	46.0	7.5	26.2	12.8	6.4	1.1			
Sunflower Oil	62.7	0.3	24.3	6.3	5.6	0.8			
^a Vegetable oils were purchased from local supermarket.									

Table S1 Distribution of fatty acids in vegetable oils^{*a*}