

ESI

Aqueous oxidation of alcohols catalysed by recoverable iron oxide
nanoparticles supported on aluminosilicates



Fatemeh Rajabi^{a*}, Antonio Pineda^b, Sareh Naserian^a, Alina Mariana Balu^{b*}, Rafael Luque^b Antonio A.
Romero^b



^aDepartment of Science, Payame Noor University, P. O. Box: 19395-4697 Tehran, Iran.



Fax: +982813344081, Tel: :+982813366366; e-mail: f_rajabi@pnu.ac.ir



^bDepartamento de Química Orgánica, Universidad de Córdoba, Campus de Rabanales, Edif. Marie Curie, Ctra Nnal IV-A, Km 396, E14014, Córdoba (Spain). Fax: +34 957212066; Tel: +34957211050;

e-mail: z82babaa@uco.es












ppm







7.8599
7.8554
7.8518
7.8423
7.8385
7.8342
7.8333
7.8292
7.8257
7.8218
7.8180
7.8142

10.0034

ppm








2.41921
2.40722

7.74715
7.73098
7.30116
7.28526


9.92618


ppm















A ¹H NMR in CDCl₃ at 289 K 90/3/1



A ¹³C NMR in CDCl₃ at 298 K 90/2/24

