

## Electronic Supplementary Information (ESI)

### Preparation of Highly Visible-light Active N-doped TiO<sub>2</sub> Photocatalyst

Guidong Yang<sup>a,b</sup>, Zheng Jiang<sup>b</sup>, Huahong Shi<sup>b</sup>, Tiancun Xiao<sup>\*b</sup>, Zifeng Yan<sup>\*\*a</sup>


<sup>a</sup> State Key Laboratory for Heavy Oil Processing, China University of Petroleum, Qingdao, 266555, China.

<sup>b</sup> Department of Chemistry, Inorganic Chemistry Laboratory, University of Oxford, OX1 3QR, U.K.


---

\*Corresponding author. Tel.: +44 1865 272660; Fax: +44 1865 272690; E-mail: [Xiao.tiancun@chem.ox.ac.uk](mailto:Xiao.tiancun@chem.ox.ac.uk).

\*\* Corresponding author. Tel.: +86 532 86981296; Fax: +86 532 86981295; E-mail address: zfyancat@upc.edu.cn



**Figure S1** MB adsorption curves on different catalysts. Adsorption conditions: 0.050 g catalyst, 50.0 mL 0.010 g/L MB, temperature = 283 K.



**Figure S2** MO adsorption curves on different catalysts. Adsorption conditions: 0.050 g catalyst, 50.0 mL 0.010 g/L MO, temperature = 283 K.