

Supporting Information

Wet Chemical Synthesis of Nitrogen-Doped Graphene towards Oxygen Reduction Electrocatalysts without High-Temperature Pyrolysis

Yuanjian Zhang,^{†*} Keisuke Fugane,[†] Toshiyuki Mori,[†] Li Niu,^{‡*} and Jinhua Ye[†]

[†]International Center for Young Scientists (ICYS), International Center for Materials
Nanoarchitectonics (MANA), National Institute for Materials Science (NIMS), 1-1
Namiki, Tsukuba 305-0044 Japan, E-mail: ZHANG.Yuanjian@nims.go.jp

[‡]Engineering Laboratory for Modern Analytical Techniques, c/o State Key Laboratory of
Electroanalytical Chemistry, Changchun Institute of Applied Chemistry, Chinese
Academy of Sciences, Renming street 5625, Changchun 130022, P. R. China; E-
mail: lniu@ciac.jl.cn




Figure S1. AFM image of graphene oxide sheets (a) and the height profile (b).




Figure S2. C1s XPS spectra of rGO (a), and rGO-N (b); N1s XPS spectra of rGO-N and rGO (c).




Figure S3. Tafel plot of HiSPECTM Pt/C(20% Pt on carbon black, Johnson Matthey) in O₂-saturated 0.1 M KOH aqueous solution.