

Supporting Information

Configurational Changes of Heme Followed of Cytochrome *c*

Folding Reaction

Jungkweon Choi^{*, a, †}, Dae Won Cho,^b Sachiko Tojo,^a Mamoru Fujitsuka,^a and Tetsuro Majima^{*, a}

^a The Institute of Scientific and Industrial Research (SANKEN), Osaka University, Mihogaoka 8-1, Ibaraki, Osaka 567-0047, Japan.

^b Department of Advanced Materials Chemistry, Korea University, Sejong Campus, Sejong 339-700, Korea.

[†] **Present Address:** Center for Nanomaterials and Chemical Reactions, Institute for Basic Science (IBS), Daejeon 305-701, Republic of Korea.




Figure S1. Steady-state Raman spectra of both ferric (a) and ferrous Cyt-c (b) in 0.5 and 4 M GdHCl solutions, respectively. ($[\text{Cyt-c}] = 50 \mu\text{M}$, $\lambda_{\text{Excitation}} = 514 \text{ nm}$).




Figure S2. Peak analysis of the ν_2 mode of 1585 cm⁻¹ for the unfolded ferric Cyt-c obtained with 416 nm excitation. The blue and red dotted lines denote the component spectra and the sum spectrum of all components, respectively.