Supporting Information

Bioactive prenylated phenyl derivatives derived from marine natural products: novel scaffold for the design of BACE inhibitors.

Javier Lopez-Ogalla¹, Esther Garcia-Palomero¹, Jorge Sanchez-Quesada¹, Laura Rubio¹, Elena Delgado¹, Pablo Garcia¹, Miguel Medina^{1,2}, Ana Castro^{1,3*}, Pilar Munoz^{1*}

 NOSCIRA S.A., Avenida de la Industria 52, 28760 Tres Cantos, Madrid, Spain
 Centro de Investigacion Biomedica en Red sobre Enfermedades Neurodegenerativas (CIBERNED), Nicolas Cabrera 1, 28049 - Madrid, Spain
 Instituto de Quimica Medica-CSIC, Juan de la Cierva 3, 28006 Madrid, Spain

List of Content

1. General information(Page 2)
2. ¹H, ¹³C and 2D NMR spectra(Page 3 to 71)

General Information:

Proton and carbon NMR spectra were obtained on 400 MHz NMR spectrometer. Chemical shifts are reported as δ values in parts per million (ppm) as referenced to residual solvent. $_1H$ NMR spectra are tabulated as follows: chemical shift, multiplicity (s = singlet, bs = broad singlet, d = doublet, t = triplet, q = quartet, m = multiplet), coupling constant(s), and number of protons.









































































































































