

SUPPLEMENTAL INFORMATION

Hijacking transferrin bound iron: protein-receptor interactions involved in iron transport in *N. gonorrhoeae*

Claire J. Parker Siburt,^{a,d} Petra L. Roulhac,^{a,d} Katherine D. Weaver,^{a,d} Jennifer M. Noto,^b Timothy A. Mietzner,^c Cynthia N. Cornelissen,^b Michael C. Fitzgerald,^{a,e} and Alvin L. Crumbliss^{a,e}

^a Department of Chemistry

Duke University

Durham, NC 27708-0346, USA

^b Department of Microbiology and Immunology

Virginia Commonwealth University

Richmond, VA 23298-0678, USA

^c Department of Molecular Genetics and Biochemistry

University of Pittsburgh School of Medicine

Pittsburgh, PA 15261, USA

^d These authors made equal contributions to this manuscript.

^e Address correspondence to alvin.crumbliss@duke.edu,

michael.c.fitzgerald@duke.edu

Supplemental Information

Figure S.1: SUPREX data obtained for apo-FbpA-Cit (open circles), apo-FbpA-Cit in the presence of TbpA⁺/B⁺ (closed circles), and FeFbpA-Cit (open squares) using an H/D exchange time of 1 hr.




Figure S.2: SUPREX data obtained for apo-FbpA-Cit (open circles), apo-FbpA-Cit in the presence of TbpA⁺/B⁺ (closed circles), apo-FbpA-Cit in the presence of TbpA⁺/B⁺ and holo-Tf (closed squares), and Fe-FbpA-Cit (open squares) using an H/D exchange time of 1 hr.

