Supporting Information for Amphiphilic Siderophore Production by Oilassociating Microbes

Michelle P. Kem, Hannah K. Zane, Stephen D. Springer, Julia M. Gauglitz, and Alison Butler*

Department of Chemistry & Biochemistry, University of California, Santa Barbara, CA 93106-9510

 Table S1. The m/z values for the 'b' ions of isolated amphibactin siderophores from ESI-MS/MS analysis.

Alcanivorax borkumensis SK2				<i>Vibrio</i> sp. S2A					<i>Vibrio</i> sp. S1B				
ʻb'	Ε	н	1	ʻb'	D	Ε	н	1	ʻb'	s	D	Ε	н
b ₃	668	670	696	b ₃	642	668	670	696	b ₃	640	642	668	670
b ₂	581	583	609	b ₂	555	581	583	609	b ₂	553	555	581	583
b ₁	409	411	437	b ₁	383	409	411	437	b ₁	381	383	409	411

Figure S1. The tandem mass spectra of amphibactins E and amphibactin I isolated from Alcanivorax borkumensis SK2. The ESI-MS/MS of amphibactin H is shown in Figure 2.





Figure S2. The tandem mass spectra of amphibactins D, E, H and I isolated from Vibrio sp. S2A.

Figure S3. The tandem mass spectra of amphibactins D, E, H and I isolated from Vibrio sp. S1B.

