

Electronic Supplementary Information (ESI)
**Enantioselective Crystallization in Miniemulsions Based on
Chiral Surfactants**

Tali Menahem and Yitzhak Mastai

Department of Chemistry, Bar-Ilan University, Ramat-Gan 52900 Israel.

E-mail: mastai@mail.biu.ac.il

Chiral surfactants preparation

Stearic acid N-hydroxy succinimide ester (SHE): Stearic acid (8.534g, 30 mmoles) was added to a solution of N-hydroxysuccinimide (3.452g, 30mmoles) in dry ethyl acetate (130ml). A solution of dicyclohexylcarbodiimide (6.19g, 30mmoles) in dry ethyl acetate was added and the reaction was left stirring overnight at room temperature. Dicyclohexylurea was removed by filtration and the filtrate was evaporated. The obtained crystals were recrystallized from ethanol yielded 8.5g (75%) of pure N-hydroxy succinimide ester of stearic acid. Stearic acid N-hydroxy succinimide ester: Melting Point 93°C. ¹H-NMR (p.p.m) (300MHz) (CDCl₃): 0.87 (3H,t), 1.27 (28H,m), 1.67 (2H,t), 2.57 (2H,t), 2.81 (4H,m). Elemental Analysis (%)H: 10.23, C: 69.29, N: 3.67.

N-Stearic (L or D) serine: A solution of stearic acid N-hydroxy succinimide ester (5g, 13mmoles) in tetrahydrofuran (130ml) was added to a solution of L or D serine (1.36g, 13mmoles) and sodium bicarbonate (1.09g, 13mmoles) in water (130ml). The reaction was left stirring overnight at temperature of 40°C. The solution was acidified to pH 2 with 1M hydrochloric acid and the tetrahydrofuran was evaporated. The product was filtered and recrystallized first from acetone and then from ethanol –water (4:1) to give 4.33g (90% yield). N-stearoyl L serine: Melting Point: 103°C. ¹H-NMR (p.p.m) (300MHz): (DMSO): 0.87 (3H,t), 1.27 (28H,m), 1.49 (2H,t), 2.12 (2H,t), 3.6 (2H,dq), 4.24 (1H,dd), 7.88 (1H,d). Elemental Analysis (%)H: 11.547, C: 66.454, N: 4.083

Figure S1

Light microscopy image chiral miniemulsions prepared at room temperature consisting 1 mg of L stearoylserine of W/O=0.03. (scale bar 50 micrometer)



