Supporting Information



Figure S1. Top) MALDI-TOF spectrum of **2**. Bottom) UV-vis spectra of **2** at a concentration of 2 mg/mL in chloroform (dashed line) and toluene (solid line).



Figure S2. SEM image of a dried sample of the fibrillar network of supramolecular polymers formed by **1** at a concentration of 20 mg/ml in toluene (scale bar represents 1000 nm).



Figure S3. NMR spectra of **1** at a concentration of 2 mg/ml in toluene-d_8; 1) 25 ºC, 2) 55 ºC, 3) 85 ºC.



Figure S4. NMR spectra of **2** at a concentration of 2 mg/ml in toluene-d_8; 1) 25 ^oC, 2) 55 ^oC, 3) 85 ^oC.



Figure S5. NMR spectra of **1** at a concentration of 2 mg/ml in $CDCl_3$; 1) 25 ºC, 2) 35 ºC, 3) 45 ºC, 4) 55 ºC.



Figure S6. NMR spectra of **2** at a concentration of 2 mg/ml in $CDCl_3$; 1) 25 ºC, 2) 35 ºC, 3) 45 ºC, 4) 55 ºC.



Figure S7. FTIR spectrum of **1** at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).



Figure S8. FTIR spectrum of **1** at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).



Figure S9. FTIR spectrum of **1** at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).



Figure S10. FTIR spectrum of 1 at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).



Figure S11. FTIR spectrum of **2** at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).



Figure S12. FTIR spectrum of **2** at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).



Figure S13. FTIR spectrum of **2** at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).



Figure S14. FTIR spectrum of **2** at a concentration of 10 mg/ml in chloroform (dashed line) and toluene (solid line).