

New Journal of Chemistry
Manuscript ID NJ-ART-09-2013-001107


Supporting information

Two Polypyridyl Copper(II) Complexes: Synthesis, Crystal
Structure and Interaction with DNA and Serum Protein in vitro


Xiao-Fei Zhao,^{ab,1} Yan Ouyang,^{a,1} Yan-Zhao Liu,^a Qiao-Juan Su,^a He Tian,^a
Cheng-Zhi Xie,^a and Jing-Yuan Xu^{a,*}

^a Tianjin Key Laboratory on Technologies Enabling Development of Clinical
Therapeutics and Diagnostics, School of Pharmacy, Tianjin Medical University,
Tianjin 300070, P. R. China

^b Department of Pharmacy, The Second Affiliated Hospital of Xing Tai Medical
College in Hebei, Xingtai 054000, P. R. China



1



2

FigS1. Packing view of complexes **1** and **2** highlighting π - π stacking and H-bonding between mononuclear units.




Fig S2. The plots of $\log[(F_0-F)/F]$ versus $\log[Q]$ of BSA at 298 K for complexes 1 and 2.




Fig.S3. Synchronous fluorescence spectra of BSA at room temperature upon addition of complexes **1** and **2**; $\Delta\lambda = 15$ nm. Both the concentrations of **1** and **2** are 0.0–6.4 M, respectively.




Fig.S4. Synchronous fluorescence spectra of BSA at room temperature upon addition of **1** and **2**; $\Delta\lambda = 60$ nm. Both the concentrations of **1** and **2** are 0.0 – 6.4 M, respectively.