

Chain of dimers to assembly of trimers. Temperature and ligand influenced formation of novel supramolecular assemblies of Cu(II) with isomeric (aminomethyl) pyridines and azide.

Sailaja S. Sunkari,^{*a} Bhagwan Kharediya,^a Satyen Saha,^b Bahjat Elrez,^{c,d}

Jean-Pascal Sutter^{*c,d}

^a Department of Chemistry, Mahila MahaVidyalaya, Banaras Hindu University, Varanasi
221 005, India

E-mail: sunkari.s7@gmail.com

Telephone: +91-9935963366, Fax Number: +91-542-2368127

^b Department of Chemistry, Faculty of Science, Banaras Hindu University, Varanasi
221 005, India

^c CNRS ; LCC (Laboratoire de Chimie de Coordination) ; 205, route de Narbonne, F-31077
Toulouse, France

^d Université de Toulouse ; UPS, INPT ; LCC ; F-31077 Toulouse, France
e-mail: sutter@lcc-toulouse.fr




Fig. S1. UV-Vis spectra of 3 in DMSO showing merging of d-d bands into a single broad band with increasing concentration.




Fig. S2. UV-Vis plot of **1** in different percentage of DMSO as solvents. The insignificant change in spectra shows the non-influential role of DMSO.