

Supporting Information

SYNTHESIS AND CHARACTERIZATIONS OF NEW TRIPHENYLAMINO-1,8-NAPHTHALIMIDES FOR ORGANIC LIGHT-EMITTING DIODE APPLICATION

Rungthiwa Arunchai,^a Taweesak Sudyoadsuk,^b Narid Prachumrak,^b Supawadee

Namuangruk,^c Vinich Promarak,^d Mongkol Sukwattanasinitt^a and Paitoon Rashatasakhon,^{a*}

^a *Organic Synthesis Research Unit, Department of Chemistry, Faculty of Science,*

Chulalongkorn University, Bangkok, 10330, Thailand

^b *Department of Chemistry and Center of Excellence for Innovation in Chemistry, Faculty of*

Science, Ubon Ratchathani University, Ubon Ratchathani, 34190, Thailand

^c *Nanoscale Simulation Laboratory, National Nanotechnology Center (NANOTEC), National*

Science and Technology Development Agency (NSTDA), Klong Luang, Pathumthani, 12120,

Thailand

^d *School of Chemistry and Center of Excellence for Innovation in Chemistry, Institute of*

Science, Suranaree University of Technology, Muang District, Nakorn Ratchasima, 30000,

Thailand

*E-mail: paitoon.r@chula.ac.th

Tel: 02-218-7620




Fig. S1 ¹H-NMR of A in CDCl₃




Fig. S2 ^{13}C -NMR of A in CDCl_3




Fig. S3 ¹H-NMR of TPN1 in CDCl₃




Fig. S4 ^{13}C -NMR of TPN1 in CDCl_3




Fig. S5 ¹H-NMR of TPN2 in CDCl₃




Fig. S6 ¹³C-NMR of TPN2 in CDCl₃




Fig. S7 ¹H-NMR of TPN3 in CDCl₃




Fig. S8 ¹³C-NMR of TPN3 in CDCl₃