

Supplementary Information Available

Conjugated Polyelectrolyte-Cisplatin Complex Nanoparticles for Simultaneous *In vivo* Imaging and Drug Tracking

Dan Ding,^a Kai Li,^a Zhenshu Zhu,^c Kan-yi Pu,^a Yong Hu,^b Xiqun Jiang,^c and Bin Liu^{*a}

^a Department of Chemical and Biomolecular Engineering, 4 Engineering Drive 4, National University of Singapore, Singapore 117576, cheliub@nus.edu.sg

^b National Laboratory of Solid State Microstructure, Department of Material Science and Engineering, Nanjing University, Nanjing, 210093, P. R. China.

^c Laboratory of Mesoscopic Chemistry and Department of Polymer Science and Engineering, College of Chemistry and Chemical Engineering, Nanjing University, Nanjing, 210093, P. R. China.

Figures




Figure S1. Size distributions of (A) CPE-g-PEG and (B) CPE-PEG-Pt nanoparticles in water.




Figure S2. 3D CLSM image of the HepG2 cancer cells after incubation with CPE-PEG-Pt nanoparticles for 2 h at 37 °C.




Figure S3. CLSM image of the HepG2 cancer cells without CPE-PEG-Pt nanoparticles incubation.




Figure S4. CLSM images of liver slices from mice intravenously injected with saline. (A) Fluorescence image; (B) Optical image.