

Supporting Information

Coupled Plasmons Induce Broadband Circular Dichroism in Patternable Films of Silver Nanoparticles with Chiral Ligands

Xavier Vidal^{a,b}, Won Jin Kim^a, Alexander Baev^a, Valentyna Tokar^{a,c}, Hongsub Jee^a, Mark. T. Swihart^{a,d} and Paras N. Prasad^{*a,e,f,g}

^a Institute for Lasers, Photonics and Biophotonics, University at Buffalo, The State University of New York, Buffalo, New York, 14260, USA.

^b Department of Physics and Astronomy, Macquarie University, New South Wales, 2109, Australia. Fax: (+61)298508115; Tel: (+61)298506368; E-mail: xavier.vidal@mq.edu.au


^c Faculty of Physics, Taras Shevchenko National University of Kyiv, 64 Volodymyrska st., Kyiv, 01601, Ukraine.

^d Department of Chemical and Biological, Engineering, University at Buffalo, The State University of New York, Buffalo, New York, 14260, USA. E-mail: swihart@buffalo.edu

^e Department of Chemistry, University at Buffalo, State University of New York, Buffalo, New York, 14260, USA. E-mail: pnprasad@buffalo.edu

^f Department of Electrical Engineering, University at Buffalo, State University of New York, Buffalo, New York, 14260, USA.

^g Department of Chemistry, Korea University.



S-Figure 1 Absorption spectra of protected *t*-BOC Ag NP film (red solid line) before patterning and deprotected *t*-BOC Ag NP film (blue dashed line) after patterning.