

Supporting information

Multifunctional superparamagnetic nanoshells: combining two-photon luminescence imaging, surface-enhanced Raman scattering and magnetic separation

Xiulong Jin,[†] Haiyan Li,[†] Shanshan Wang,[†] Ni Kong,[†] Hong Xu,[†] Qihua Fu,[‡]

Hongchen Gu,[†] and Jian Ye^{†,}*

[†]Shanghai Engineering Research Center of Medical Device and Technology at Med-X,
School of Biomedical Engineering, Shanghai Jiao Tong University, 1954 Huashan
Road, Shanghai, 200030, China;

[‡]Department of Laboratory Medicine, Shanghai Children's Medical Center & School
of Medicine, Shanghai Jiao Tong University, Shanghai, 200127, China

* To whom correspondence should be addressed. E-mail: yejian78@sjtu.edu.cn




Figure S1. Dependence plot of TPL intensity on the pulse laser excitation wavelength. Red line is only for visual guidance.




Figure S2. Transmission images of 4-MBA labeled $\text{Fe}_3\text{O}_4@\text{SiO}_2@\text{Au}$ nanoparticle chains self-assembled under an external magnetic field along different directions. The white double-arrow symbol in each panel indicates the direction of the magnetic field. All scale bars are $50\ \mu\text{m}$.