Electronic Supplementary Material (ESI) for Nanoscale

Horseradish peroxidase-immobilized magnetic mesporous silica

nanoparticles as a potential candidate to eliminate intracellular

reactive oxygen species

YajingShen^a, Ye Zhang^a, Xiang Zhang^a, XiuhongZhou^b, XiyaoTeng^a,

Manqing Yan^a, and Hong Bi*,a

^aCollege of Chemistry and Chemical Engineering, Anhui University, Hefei 230601, China

^bBiotechnology Center of Anhui Agricultural University, Hefei 230601, China

*Corresponding author: Fax/Tel: +86 551 63861279; E-mail: <u>bihong@ahu.edu.cn</u>

Supporting Information

Table of content

Fig. S1 TEM image of CDs (The inset shows the histogram of size distribution)

Fig. S2 XRD patterns of (a) Fe₃O₄ nanoparticles and (b) MMSNs.

- Fig. S3 N₂ adsorption/desorption isotherms of MMSNs (the inset is the pore size distribution histogram).
- Fig. S4 FT-IR spectra of (a) Fe₃O₄ nanoparticles, (b) MMSNs and (c) MMSNs-NH₂.
- Fig. S5 PL emission spectra of 0.5 mg CDs dispersed in aqueous solution containing different concentrations of (a) HRP and (b) MMSNs-HRP ($\lambda_{ex} = 365$ nm).
- Fig. S6 Optical microscopic images of CHO cells treated with different concentrations of H_2O_2 (600 μ M, 800 μ M, 1000 μ M, and 2000 μ M) for 20 h.
- Fig. S7 The fluorescent microscopy images of CHO cells co-incubated with MMSNs-HRP at different concentrations of 100, 200, 300 and 400 μ g mL⁻¹ for 44 h.



Fig. S1 TEM image of CDs (The inset shows the histogram of size distribution)



Fig. S2 XRD patterns of (a) Fe₃O₄ nanoparticles and (b) MMSNs.



Fig. S3 N_2 adsorption/desorption isotherms of MMSNs (the inset is the pore size distribution curve).



Fig. S4 FT-IR spectra of (a) Fe_3O_4 nanoparticles, (b) MMSNs and (c) MMSNs-NH₂.



Fig. S5 PL emission spectra of 0.5 mg CDs dispersed in aqueous solution containing different concentrations of (a) HRP and (b) MMSNs-HRP ($\lambda_{ex} = 365$ nm).



Fig. S6 Optical microscopic images of CHO cells treated with different concentrations of H_2O_2 (600 μ M, 800 μ M, 1000 μ M, and 2000 μ M) for 20 h.

100 μg/mL	a	200 μg/mL	b	300 μg/mL	c	400 μg/mL	d
	9						

Fig. S7 The fluorescent microscopy images of CHO cells co-incubated with MMSNs-HRP at different concentrations of 100, 200, 300 and 400 μ g mL⁻¹ for 44 h.