Biosynthetic Pathways to Isocyanides and Isothiocyanates; Precursor Incorporation Studies on Terpene Metabolites in the Tropical Marine Sponges *Amphimedon terpenensis* and *Axinyssa* n.sp.

Jamie S. Simpson, and Mary J. Garson

Supplementary data

Table of Contents

Figure S1. ¹ H NMR spectrum for formamide 14	S2
Figure S2. ¹³ C NMR spectrum for formamide 14	S2
Figure S3. ¹ H NMR spectrum for thiourea 15	S3
Figure S4. ¹³ C NMR spectrum for thiourea 15	S3
Figure S5. ¹ H NMR spectrum for thiol 16	S4
Figure S6. ¹³ C NMR spectrum for thiol 16	S4
Figure S7. ¹ H NMR spectrum for bisisothiocyanate 17	S5
Figure S8. ¹³ C NMR spectrum for bisisothiocyanate 17	S5
Figure S9. ¹ H NMR spectrum spectra for amine 18	S 6
Figure S10. ¹³ C NMR spectrum for amine 18	S6



32



Figure S3. ¹H NMR spectrum for thiourea 15

160

140

120

100

80

40

Supplementary Material for Organic & Biomolecular Chemistry



Figure S7. ¹H NMR spectrum for bisisothiocyanate 17



