

Synthesis and characterisation of estrogenic carriers for cytotoxic Pt(II) fragments: biological activity of the resulting complexes

Supporting Information

Elisabetta Gabano,^a Claudio Cassino,^a Samuele Bonetti,^a Cristina Prandi,^a Donato Colangelo,^b AnnaLisa Ghiglia^b and Domenico Osella^a*

^a *Dipartimento di Scienze dell'Ambiente e della Vita, Università del Piemonte Orientale "A. Avogadro", Spalto Marengo 33, 15100 Alessandria, Italy. Fax: +39 0131 287416; Tel: +39 0131 287429; E-mail: domenico.osella@mfn.unipmn.it*

^b *Dipartimento di Scienze Mediche, Università del Piemonte Orientale "A. Avogadro", via Solaroli 17, I-28100, Novara, Italy.*




Figure S1. ^1H -NMR spectrum of ligand 4 in D_2O .




Figure S2. ¹³C-NMR spectrum of ligand 4 in D₂O.




Figure S3. $^1\text{H-NMR}$ spectrum of complex 7 in $(\text{CD}_3)_2\text{SO}$.




Figure S4. ^{13}C -NMR spectrum of complex 7 in $(\text{CD}_3)_2\text{SO}$.




Figure S5. ^{195}Pt -NMR spectrum of complex 7 in $(\text{CD}_3)_2\text{SO}$.




Figure S6. $^1\text{H-NMR}$ spectrum of ligand **6** in $(\text{CD}_3)_2\text{CO}$.




Figure S7. ^{13}C -NMR spectrum of ligand **6** in $(\text{CD}_3)_2\text{CO}$.




Figure S8. ¹H-NMR spectrum of complex 8 in (CD₃)₂SO.




Figure S9. ^{13}C -NMR spectrum of complex 8 in $(\text{CD}_3)_2\text{SO}$.




Figure S10. ^{195}Pt -NMR spectrum of complex 8 in $(\text{CD}_3)_2\text{SO}$.