

Electronic Supplementary Information

Intramolecular Direct Arylation in an A,C-functionalized Calix[4]arene

Olaf G. Barton,^a B. Neumann,^b H.-G. Stammer^b and Jochen Mattay^{*a}

^a*Organische Chemie I. Fax: +49 521 106 6417; Tel: +49 521 106 2072;*

E-mail: ocljm@uni-bielefeld.de

^b*Abteilung für Röntgenstrukturanalyse, Anorganische Chemie II.*

^{a,b}*Universität Bielefeld, Universitätsstr. 25, 33501 Bielefeld, Germany.*

Table of Contents

NMR spectra of 4	S1
NMR spectra of 2	S2
Variable temperature ^1H NMR spectra 1a	S3
Variable temperature ^1H NMR spectra 1b	S4
^{13}C NMR spectrum of 1a	S5
^{13}C NMR spectrum of 1b	S6
NMR spectra of 8	S7
NMR spectra of 6	S8




Fig. S1 ^1H and ^{13}C NMR spectrum of **4** in CHCl_3 .




Fig. S2 ^1H and ^{13}C NMR spectrum of **2** in CHCl_3 .




Fig. S3 Variable temperature ^1H NMR spectra **1a**, residual solvent signals are marked (*).




Fig. S4 Variable temperature ^1H NMR spectra **1b**, residual solvent signals are marked (*).




Fig. S5 ¹³C NMR spectrum of **1a** in C₂D₂Cl₄.




Fig. S6 ¹³C NMR spectrum of **1b** in CD₂Cl₂.




Fig. S7 ^1H and ^{13}C NMR spectrum of **8** in CHCl_3 .




Fig. S8 ^1H and ^{13}C NMR spectrum of **6** in CHCl_3 .