

Supplementary Data

Cyclic triureas – synthesis, crystal structures and properties

Denys Meshcheryakov,^a Françoise Arnaud-Neu,^b Volker Böhmer,^{*a} Michael Bolte,^c Véronique Hubscher-Bruder,^b Emilie Jobin,^b Iris Thondorf^d and Sabine Werner^d

^aAbteilung Lehramt Chemie, Fachbereich Chemie, Pharmazie und Geowissenschaften, Johannes Gutenberg-Universität Mainz, Duesbergweg 10-14, D-55099, Mainz, Germany, ^bIPHC-DSA, ULP, CNRS ; 25 rue Becquerel, F-67087 Strasbourg, France, ^cFachbereich Chemie und Pharmazeutische Wissenschaften, Institut für Anorganische Chemie, Johann Wolfgang Goethe-Universität Frankfurt, Marie-Curie-Straße 11 , D-60439 Frankfurt/Main, Germany and ^dInstitut für Biochemie und Biotechnologie, Fakultät für Naturwissenschaften I Biowissenschaften, Martin-Luther-Universität Halle-Wittenberg, Kurt-Mothes-Str. 3, D-06099 Halle, Germany

Contents

Table S1. Compounds used for the calculation of RESP charges and the corresponding RESP derived charges for the constituting fragments of them macrocycles **10-13**. Also given are the corresponding atom types used for the AMBER calculations.

Table. S1 Compounds used for the calculation of RESP charges and the corresponding RESP derived charges for the constituting fragments of them macrocycles **10-13**. Also given are the corresponding atom types used for the AMBER calculations.

Molecule for the RESP calculation	Fragment in 10 - 13