

Supplementary materials

Selectivity and Affinity of DNA Triplex Forming Oligonucleotides Containing the Nucleoside Analogues 2'-*O*-Methyl-5-(3-amino-1-propynyl)uridine and 2'-*O*-Methyl-5-propynyluridine

Hong Li,^a Paul S. Miller,^b and Michael M. Seidman*^a

^a*Laboratory of Molecular Gerontology, National Institute on Aging, National Institutes of Health, Baltimore, Maryland 21224. E-mail: seidmanm@grc.nia.nih.gov; fax :+01- 410-558-8157; Tel : + 01-410-558-8565*

^b*Department of Biochemistry and Molecular Biology, Bloomberg School of Public Health, Johns Hopkins University, Baltimore, Maryland 21205. E-mail: pmiller@jhsph.edu; fax :+01- 410-955-2926; Tel : + 01- 410-955-3489*

Table S1 Maldi-TOF data of Oligos

Oligonucleotides	(M+H) ⁺ calculated	(M+H) ⁺ found
5' -TTTCTCTTTTCTTCTAC-3' 5	5653	5653
3' -AAAGAGAAAAAGAAGATG-5' 6	5961	5962
3' -TTTCTCTTTTCTTCT-5' 7	5616	5617
3' -TTTCTC 1 TTTT 1 C 1 TC 1 -5' 8	5712	5713
3' -TTTCTCT 1 T 1 T 1 C 1 TCT-5' 9	5712	5712
3' -TTTCTCT 1 1 1 TCTTCT-5' 10	5712	5713
3' -TTTCTC 2 TTTT 2 C 2 TC 2 -5' 11	5772	5772
3' -TTTCTCT 2 T 2 T 2 C 2 TCT-5' 12	5772	5772
3' -TTTCTCT 2 2 2 TCTTCT-5' 13	5772	5773
3' - T4434 CTTTTTCTTCT-5' 14	5732	5733
3' - T4434 CT 2 T 2 T 2 C 2 TCT-5' 15	5888	5888
3' - T4434 C 2 TTTC 2 CT-5' 16	5888	5889
3' - T4434 C 2 T 2 TTTCT 2 C 2 -5' 17	5888	5889

In oligos **7-17**, T is 2'-O-Me-5-metheyl dU, C is 2'-O-Me-5-methyl dC. **1**: 2'-O-Methyl-5-propynyluridine, **2**: 2'-O-Methyl-5-(3-amino-1-propynyl)uridine, **3**: 2'-aminoethoxy-5-metheyl dC, **4**: 2'-aminoethoxy-5-metheyl dU.

Figure S1




Fig. S1 The profile of non-denaturing gel electrophoresis of triplexes.