

Supplementary Information:

Asymmetric transfer hydrogenation of imines catalyzed by polymer-immobilized chiral catalyst

Naoki Haraguchi*, Keiichi Tsuru, Yukihiro Arakawa, and Shinichi Itsuno

^a 1-1 Hibarigaoka, Tempaku-cho, Toyohashi, Aichi 441-8580, Japan.
E-mail: haraguchi@tutms.tut.ac.jp; Tel: +81 532 44 6812; Fax: +81 532 48 5833

Table of Contents

¹H and ¹³C NMR

Figure S1. ¹ H-NMR spectrum of (1 <i>R</i> ,2 <i>R</i>)- <i>N</i> ¹ -(4-vinylbenzenesulfonyl)-1,2-diphenylethane-1,2-diamine.	S2
Figure S2. ¹ H-NMR spectrum of 4-vinylbenzenesulfonic acid benzyltributylammonium salt.	S2
Figure S3. ¹ H-NMR spectrum of (<i>R,R</i>)- 1 .	S3
Figure S4. ¹ H-NMR spectrum of (<i>R,R</i>)- 3 .	S4
Figure S5. ¹³ C-NMR spectrum of (<i>R,R</i>)- 3 .	S4
Figure S6. ¹ H-NMR spectrum of (<i>R</i>)- 5 .	S5
Figure S7. ¹ H-NMR spectrum of (<i>S</i>)- 11 .	S5




Figure S1. ¹H-NMR spectrum of (1*R*,2*R*)-*N*¹-(4-vinylbenzenesulfonyl)-1,2-diphenylethane-1,2-diamine.




Figure S2. ¹H-NMR spectrum of 4-vinylbenzenesulfonic acid benzyltributylammonium salt.




Figure S3. ^1H -NMR spectrum of (R,R) -1.




Figure S4. ^1H -NMR spectrum of (R,R) -3.




Figure S5. ^{13}C -NMR spectrum of (R,R) -3.




Figure S6. ¹H-NMR spectrum of (*R*)-5.




Figure S7. ¹H-NMR spectrum of (*S*)-11.