



Electronically Tuned 1,3,5-Triarylpyrazolines as Cu(I)-Selective Fluorescent Probes


Manjusha Verma, Aneese F. Chaudhry, M. Thomas Morgan, and Christoph J. Fahrni

Supporting Information


Table of Contents



1. ¹ H-NMR spectra of compounds 4 and 5	S2
2. ¹ H-NMR and ¹⁹ F-NMR spectra of compound 6	S3
3. ¹ H-NMR and ¹⁹ F-NMR spectra of compound (±)- 1a	S4
4. ¹ H-NMR and ¹⁹ F-NMR spectra of compound (±)- 1b	S5
5. ¹ H-NMR and ¹⁹ F-NMR spectra of compound (±)- 1c	S6
6. ¹ H-NMR and ¹⁹ F-NMR spectra of compound (±)- 1d	S7
7. ¹ H-NMR and ¹⁹ F-NMR spectra of compound (±)- 1e	S8
8. ¹ H-NMR and ¹⁹ F-NMR spectra of compound (±)- 1f	S9






¹⁹F-NMR (CDCl₃, 376 MHz)








¹H-NMR (CDCl₃, 400 MHz)


Supplementary Material (ESI) for Organic & Biomolecular Chemistry
This journal is (c) The Royal Society of Chemistry 2010




¹⁹F-NMR (CDCl₃, 376 MHz)










¹⁹F-NMR (CDCl₃, 376 MHz)







¹⁹F-NMR (CDCl₃, 376 MHz)

