

Electronic Supplementary Information for

A combined spin trapping / EPR / mass spectrometry approach to study the formation of a cyclic peroxide by dienolic precursor autoxidation

Mathilde Triquigneaux,^a Laurence Charles,^a Christiane André-Barrès^b and Béatrice Tuccio*^a

^a Universités Aix-Marseille I, II & III – CNRS, UMR 6264: Laboratoire Chimie Provence, Spectrométries Appliquées à la Chimie Structurale, 13397 Marseille Cedex 20, France .
Fax: (+33) 491282897; Tel: (+33) 491288743;
E-mail: beatrice.tuccio-lauricella@univ-provence.fr

^b Université Paul Sabatier – CNRS, UMR 5068: Synthèse et Physico-Chimie de Molécules d’Intérêt Biologique, 31062 Toulouse cedex, France

Table of contents

Figure S1 : ESI-MS/MS spectra of a) $[1\mathbf{a}+\mathbf{H}]^+$ at 30 eV and b) $[1\mathbf{a}-\mathbf{H}]^-$ at 30 eV.

Figure S2 : ESI-MS/MS spectra of a) $[1\mathbf{8}+\mathbf{H}]^+$ at 5 eV and b) $[1\mathbf{8}-\mathbf{H}]^-$ at 5 eV.




Figure S1 : ESI-MS/MS spectra of a) $[1\mathbf{a}+\mathbf{H}]^+$ at 30 eV and b) $[1\mathbf{a}-\mathbf{H}]^-$ at 30 eV.




Figure S2 : ESI-MS/MS spectra of a) $[18+\text{H}]^+$ at 5 eV and b) $[18-\text{H}]^-$ at 5 eV.