

Supplementary Information

Extending conjugation in porphyrin dimer carbocations

Karl J. Thorley and Harry L. Anderson*

Department of Chemistry,
Chemistry Research Laboratory,
University of Oxford,
12 Mansfield Road,
Oxford,
UK
OX1 3TA.
E-mail: harry.anderson@chem.ox.ac.uk;
Fax: +44 1865 285002;
Tel: +44 1865 275704

Contents

NMR spectra:

[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-butadiynyl porphinato] nickel(II) 3	1
[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-hexatriynyl porphinato] nickel(II) 4	2
[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-[ethynyl- <i>p</i> -(trimethylsilylethynyl)phenyl] porphinato] nickel(II) 10	3
[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-(ethynylphenone) porphinato] nickel(II) 17	4
[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-ethynyl porphinato] [5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl porphinato] methanolato nickel(II) 13	5
[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-butadiynyl porphinato] [5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl porphinato] methanolato nickel(II) 15	7
[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-butadiynyl porphinato] [5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-ethynyl porphinato]benzenemethanolato nickel(II) 18	9
bis[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-butadiynyl porphinato] benzenemethanolato nickel(II) 6.....	10
bis[5,15-bis(3,5-di- <i>tert</i> -butylphenyl)-10-phenyl-20-[ethynyl- <i>p</i> -(trimethylsilylethynyl)phenyl] porphinato] benzenemethanolato nickel(II) 11.....	12

[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-butadiynyl porphinato] nickel(II) 3




Figure S1. ^1H NMR spectra of porphyrin monomer 3, recorded in CDCl_3 at 300 K, 400 MHz.

[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-hexatriynyl porphinato] nickel(II) 4




Figure S2. ¹H NMR spectra of porphyrin monomer 4, recorded in CDCl₃ at 300 K, 400 MHz.

[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-[ethynyl-*p*-(trimethylsilyl)ethynyl]porphinato] nickel(II) **10**




Figure S3. ¹H NMR spectra of porphyrin monomer **10**, recorded in CDCl₃ at 300 K, 400 MHz.

[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-(ethynylphenone) porphinato] nickel(II) 17




Figure S4. ¹H NMR spectra of porphyrin monomer 17, recorded in CDCl₃ at 300 K, 400 MHz.

[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-ethynyl porphinato] [5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl porphinato] methanolato nickel(II) **13**




Figure S5. ¹H NMR spectra of porphyrin dimer **13**, recorded in CDCl₃ at 300 K, 400 MHz.




Figure S6. ¹³C NMR spectrum of porphyrin dimer **13**, recorded in CDCl₃ at 300 K, 125 MHz.

[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-butadiynyl porphinato] [5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl porphinato] methanolato nickel(II) 15




Figure S7. ¹H NMR spectra of porphyrin dimer 15, recorded in CDCl₃ at 300 K, 400 MHz.




Figure S8. ¹³C NMR spectrum of porphyrin dimer **15**, recorded in CDCl₃ at 300 K, 125 MHz.

[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-butadiynyl porphinato] [5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-ethynyl porphinato]benzenemethanolato nickel(II) **18**




Figure S9. ¹H NMR spectra of porphyrin dimer **18**, recorded in CDCl₃ at 300 K, 400 MHz.

bis[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-butadiynyl porphinato] benzenemethanolato nickel(II) 6




Figure S10. ¹H NMR spectra of porphyrin dimer 6, recorded in CDCl₃ at 300 K, 400 MHz.




Figure S11. ¹³C NMR spectrum of porphyrin dimer **6**, recorded in CDCl₃ at 300 K, 125 MHz.

bis[5,15-bis(3,5-di-*tert*-butylphenyl)-10-phenyl-20-[ethynyl-*p*-(trimethylsilylethynyl)phenyl]porphinato] benzenemethanolato nickel(II) 11




Figure S12. ¹H NMR spectra of porphyrin dimer 11, recorded in CDCl₃ at 300 K, 400 MHz.