

Supporting Information for Org. Biomol. Chem.

Synthesis of chiral calix[4]arenes bearing aminonaphthol moieties and
their use in the enantiomeric recognition of carboxylic acids

Mustafa Durmaz, Mustafa Yilmaz and Abdulkadir Sirit*

*Department of Chemistry, Selçuk University, 42099 Konya, Turkey
E-mail: asirit42@yahoo.com*

Table of Contents

S2.	¹ H NMR spectra of 1
S2.	¹ H NMR spectra of 2
S3.	¹ H NMR spectra of 3
S3.	¹ H NMR spectra of 4
S4.	¹ H NMR spectra of 8
S4.	¹³ C NMR spectra of 8
S5.	¹ H NMR spectra of 9
S5.	¹³ C NMR spectra of 9
S6.	¹ H NMR spectra of 10
S6.	¹³ C NMR spectra of 10
S7.	¹³ C DEPT NMR spectra of 10 (CDCl ₃)
S8.	¹ H NMR spectra of 11
S8.	¹³ C NMR spectra of 11
S9.	¹ H NMR spectra of 12
S9.	¹³ C NMR spectra of 12
S10.	¹ H NMR spectra of 13
S10.	¹³ C NMR spectra of 13
S11.	¹ H NMR spectra of 14
S11.	¹³ C NMR spectra of 14
S12.	¹ H NMR spectra of 15
S12.	¹³ C NMR spectra of 15
S13.	¹ H NMR spectra of 16
S13.	¹³ C NMR spectra of 16
S14.	GCOSY NMR spectra of 16 (CDCl ₃)
S15.	¹ H NMR titration curves of compound 9 with (<i>R</i>)- and (<i>S</i>)- mandelic acids
S15.	Job plot analysis of compound 16 with (<i>R</i>)- and (<i>S</i>)- mandelic acids
S15.	Typical Benesi-Hildebrand plot of 1/ΔA versus 1/[G] ₀ (left) and the plot of ln K versus 1/T for the host–guest complexation of 16 and (<i>D</i>)-DBTA in CHCl ₃ at 25°C (right)
S15.	UV/Vis spectra of chiral calix[4]arene 16 (1.00 × 10 ⁻⁵ M) in the presence of (<i>D</i>)-Dibenzoyltartaric acid (0.10–3.00×10 ⁻⁴ M) in CHCl ₃ .



























