



Electronic Supplementary Information



Index


Contents of Supporting Information page	1
General Experimental Protocols	2
NMR Spectra	
2-Propylsulfanyl-benzothiazole 1	3
2-(Propane-1-sulfonyl)-benzothiazole 2	4
1-Phenyl-5-propylsulfanyl-1 <i>H</i> -tetrazole 3	5
1-Phenyl-5-(propane-1-sulfonyl)-1 <i>H</i> -tetrazole 4	6
1-Methyl-4-(propane-1-sulfonyl)-benzene 5	7
2-Bromoethyl benzene 6	8
<i>n</i> -Propyl triphenylphosphonium bromide 7	9
<i>tert</i> -Butyl-(2,2-dichloro-11-iodo-undecyloxy)-dimethyl-silane 8	10
References	11

General Experimental Protocols


2-Propylsulfanyl-benzothiazole, 2-(propane-1-sulfonyl)-benzothiazole, 1-phenyl-5-propylsulfanyl-1*H*-tetrazole, 1-phenyl-5-(propane-1-sulfonyl)-1*H*-tetrazole, 1-methyl-4-(propane-1-sulfonyl)-benzene and propyl triphenylphosphonium bromide were prepared by the literature methods.^{31, 40, 45} All proton nmr spectra were recorded in deuteriochloroform and referenced relative to residual chloroform at δ 7.26. Spectra were recorded using 32k data points with a sweep width of 8224 Hz, and processed with one level of zero filling with a line broadening of 0.3 Hz giving a digital resolution of 0.25 Hz. Spectra were simulated and iteratively analysed using the gNMR v3.6 for Windows software package.











3









4









References

31. A.K. Ghosh, B. Zajc, *J. Org. Chem.*, **2009**, *74*, 8531-8540.
40. P. Oxley, M.W. Partridge, D.T. Robson, W.F. Short, *J. Chem. Soc.*, **1946**, 763-771.
45. J. Bourgeois, I. Dion, P.H. Cebrowski, F. Loiseau, A.C. Bedard, A.M. Beauchemin, *J. Am. Chem. Soc.*, **2009**, *131*, 874-875.