

## *Supporting Information*

# **Syntheses of sulfur and selenium analogues of pachastrissamine via double displacements of cyclic sulfate**

**Hongjun Jeon<sup>a</sup>, Hoon Bae<sup>a</sup>, Dong Jae Baek<sup>a</sup>, Young-Shin Kwak<sup>b</sup>, Deukjoon Kim<sup>a</sup>,  
and Sanghee Kim<sup>a,\*</sup>**

<sup>a</sup> College of Pharmacy, Seoul National University, San 56-1, Shilim, Kwanak, Seoul  
151-742, Korea

<sup>b</sup> Korea Research Institute of Bioscience & Biotechnology (KRIBB), Chungbuk 363-  
883, Korea

*pennkim@snu.ac.kr*

## **Contents**

<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>7</b>	S3-S4
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>8</b>	S5-S6
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>9</b>	S7-S8
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>10</b>	S9-S10
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>11</b>	S11-S12
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>12</b>	S13-S14
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>15</b>	S15-S16
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>16</b>	S17-S18
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>17</b>	S19-S20
<sup>1</sup> H NMR and <sup>13</sup> C NMR spectra of compound <b>18</b>	S21-S22

$^1\text{H}$ NMR and $^{13}\text{C}$ NMR spectra of compound <b>4</b>	S23-S24
$^1\text{H}$ NMR and $^{13}\text{C}$ NMR spectra of compound <b>5</b>	S25-S26





















































