

Supplementary Material

An efficient approach to dispacamide A and its derivatives.

Solène Guihéneuf, Ludovic Paquin, François Carreaux, Emilie Durieu,
Laurent Meijer and Jean Pierre Bazureau


*Université de Rennes 1, Laboratoire Sciences Chimiques de Rennes,
UMR CNRS 6226, Groupe Ingénierie Chimique & Molécules pour le Vivant (ICMV),
Bât. 10A, Campus de Beaulieu, Avenue du Général Leclerc, CS 74205
35042 Rennes Cedex, France.*

*Fax: +33 (0)223 236 374. Phone: +33(0)2 223 236 603.
E-mail: jean-pierre.bazureau@univ-rennes1.fr*


ELECTRONIC SUPPLEMENTARY INFORMATION

¹³C NMR spectra of isolated compounds 11(a-f)	p2-p7
¹H NMR spectra of isolated compounds 11(a-f)	p8-p13


¹³C NMR spectrum of dispacamide A (11a)




^{13}C NMR spectrum of compound (11b)




¹³C NMR spectrum of compound (11c)




¹³C NMR spectrum of compound (11d)




^{13}C NMR spectrum of compound (11e)




^{13}C NMR spectrum of compound (11f)




¹H NMR spectrum of dispacamide A (11a)




¹H NMR spectrum of compound (11b)



¹H NMR spectrum of compound (11c)



¹H NMR spectrum of compound (11d)



¹H NMR spectrum of compound (11f)

