## Supplementary Information

## A smooth rearrangement of N-p-toluenesulfonyl 2-tert-butyldiphenylsilylmethyl-substituted azetidines into N-p-toluenesulfonyl 3-tert-butyldiphenylsilyl-substituted pyrrolidines

Bharat D. Narhe, <sup>a</sup> Vardhineedi Sriramurthy, <sup>b</sup> Veejendra K. Yadav <sup>a</sup>\*

<sup>a</sup>Department of Chemistry, Indian Institute of Technology, Kanpur 208 016, India.

<sup>b</sup>Current address: Chemical Development, TCI America, 9211 N. Harborgate Street, Portland, OR 97203, USA

.....

| S1-S3      | Table of contents   |
|------------|---|
| S4         | $^{1}$ H and $^{13}$ C NMR spectra of (S)- $p$ -toluenesulfinylimine of 2- $tert$ -butyldiphenylsilylacetaldehyde |
| <b>S</b> 5 | $^{1}$ H and $^{13}$ C NMR spectra of <b>5 (R = H)</b>  |
| <b>S</b> 6 | $^{1}$ H and $^{13}$ C NMR spectra of <b>5a (R = Me)</b>  |
| <b>S7</b>  | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>5b (R = Me)</b>  |
| <b>S8</b>  | $^{1}$ H and $^{13}$ C NMR spectra of <b>5a</b> ( <b>R</b> = allyl)   |
| <b>S</b> 9 | $^{1}$ H and $^{13}$ C NMR spectra of <b>5b</b> ( <b>R</b> = allyl)   |
| S10        | $^{1}$ H and $^{13}$ C NMR spectra of <b>5a</b> ( <b>R</b> = <b>benzyl</b> )                                      |
| S11        | $^{1}$ H and $^{13}$ C NMR spectra of <b>5b</b> ( <b>R</b> = <b>benzyl</b> )                                      |
| S12        | $^{1}$ H and $^{13}$ C NMR spectra of <b>5a</b> ( <b>R</b> = <b>vinyl</b> )                                       |
| S13        | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>5b (R = vinyl)</b>   |
| S14        | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>5a (R = phenyl)</b>  |
| S15        | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>5b (R = phenyl)</b>  |
| S16        | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>5a (R = <i>p</i>-fluorophenyl)</b>                           |
| S17        | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>5b</b> ( <b>R</b> = $p$ -fluorophenyl)                       |
| S18        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7 (R = H)</b>  |
| S19        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7a</b> ( <b>R = Me</b> )   |
| S20        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7b</b> ( <b>R</b> = <b>Me</b> )  |
| S21        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7a</b> ( <b>R</b> = allyl)   |
| S22        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7b</b> ( <b>R</b> = allyl)   |
| S23        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7a</b> ( <b>R</b> = <b>benzyl</b> )                                      |
| S24        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7b</b> ( <b>R</b> = <b>benzyl</b> )                                      |
| S25        | $^{1}$ H and $^{13}$ C NMR spectra of <b>7a</b> ( <b>R</b> = <b>vinyl</b> )                                       |
| S26        | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>7b (R = vinyl)</b>   |
| S27        | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>7a (R = phenyl)</b>  |
|            |   |

```
<sup>1</sup>H and <sup>13</sup>C NMR spectra of 7b (R = phenyl)
S28
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 7a (R = p-fluorophenyl)
S29
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 7b (R = p-fluorophenyl)
S30
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of N-p-Toluenesulfonyl-2(R)-tert-butyldiphenylsilylmethylazetidine
S31
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1b<sub>1</sub>
S32
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1b<sub>2</sub>
S33
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1f<sub>1</sub>
S34
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1f<sub>2</sub>
S35
S36
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1c<sub>1</sub>
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1c<sub>2</sub>
S37
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1g<sub>1</sub>
S38
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1g<sub>2</sub>
S39
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1d<sub>1</sub>
S40
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1d<sub>2</sub>
S41
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1e<sub>1</sub>
S42
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 1e<sub>2</sub>
S43
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of (±)-3a
S44
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3b<sub>1</sub>
S45
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3b<sub>2</sub>
S46
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of (E)-13
S47
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3f<sub>1</sub>
S48
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3f<sub>2</sub>
S49
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of (E/Z)-16
S50
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3c<sub>1</sub>
S51
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3c<sub>2</sub>
S52
                 ^{1}H and ^{13}C NMR spectra of (E/Z)-14
S53
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3g<sub>1a</sub>
S54
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of (±)-3g
S55
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3g<sub>2a</sub>
S56
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3d<sub>1</sub>
S57
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of (E)-15d
S58
S59
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3d<sub>2</sub>
                 <sup>1</sup>H and <sup>13</sup>C NMR spectra of 3e<sub>1</sub>
S60
```

| S61 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of ( <i>E</i> )- <b>15e</b>       |
|-----|--|
| S62 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>3e<sub>2</sub></b>          |
| S63 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of (±)- <b>19a</b>                |
| S64 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>20b<sub>2</sub></b>         |
| S65 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>10</b>                      |
| S66 | $^{1}\text{H}$ and $^{13}\text{C}$ NMR spectra of 12 ( $\alpha\text{-OH})$       |
| S67 | $^{1}\text{H}$ and $^{13}\text{C}$ NMR spectra of <b>12</b> ( $\beta\text{-OH})$ |
| S68 | $^{1}$ H and $^{13}$ C NMR spectra of $\mathbf{1h_{1}}$                          |
| S69 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>1h<sub>2</sub></b>          |
| S70 | $^{1}$ H and $^{13}$ C NMR spectra of $\mathbf{3h_{1}}$                          |
| S71 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>17</b>                      |
| S72 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>3h<sub>2</sub></b>          |
| S73 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>18</b>                      |
| S74 | $^{1}$ H and $^{13}$ C NMR spectra of $\mathbf{21h_{1}}$                         |
| S75 | <sup>1</sup> H and <sup>13</sup> C NMR spectra of <b>21h<sub>2</sub></b>         |
|     |  |



























































































































































































































































