

Supplementary Information

Non-peptidic cell-penetrating agents: Synthesis of oligomeric chiral bicyclic guanidinium vectors

Julián Valero,^a Michiel Van Gool,^b Ruth Pérez-Fernández,^{a,d} Pilar Castreño,^a Jorge Sánchez-Quesada,^b Pilar Prados,^b and Javier de Mendoza^{*,a,c}

^aInstitute of Chemical Research of Catalonia (ICIQ), Avda. Països Catalans 16, E-43007 Tarragona, Spain; ^bDepartamento de Química Orgánica, Facultad de Ciencias, Universidad Autónoma de Madrid, Cantoblanco, E-28049 Madrid, Spain;

^cDepartament de Química Analítica i Química Orgànica, Facultat de químiques, Universitat Rovira i Virgili, Campus de Sescelades, E-43007 Tarragona, Spain and ^dCurrent address: Instituto de Química Médica, CSIC, Juan de la Cierva, 3, E-28006 Madrid, Spain.

*To whom correspondence should be addressed. Phone: 34 977 920 220. Fax: 34 977 920 226. E-mail: jmendoza@iciq.es


Table of contents:

Preparation of PF ₆ ⁻ impregnated silica	S2
Relevant NMR spectra	S3-S32
HPLC profiles of Compounds 9 , 10 , 11 , 12 , 15 , 20 , 21 , 22 and 29	S33-S39


Preparation of silica with KPF_6^-

To silica gel (40-60 μm , 100 g), KPF_6 (3 g) and 200 mL of acetone were added. The mixture was stirred for two hours and the solvent was evaporated under vacuum. The resulting solid was placed overnight into an oven (50-70°C).


Compound 2




Compound 4




Compound 5




Compound 7




Compound 8




Compound 9





Compound 10






Compound 11




Compound 13




Compound 14




Compound 16




Compound 17




Compound 18





Compound 20





Compound 21




Compound 23




Compound 24




Compound 25




Compound 26






Compound 27





Compound 28




Compound 29




Compound 31









Compound 32





Compound 33




Compound 34




Compound 35



Compound 36





Compound 37




HPLC Profiles


Compound 9 HPLC-MS





Compound 10




Compound 11





Compound 12 HPLC-MS









Compound 15




Compound 20 HPLC-MS





Compound 21 HPLC-MS



Compound 22 HPLC-MS



22



Compound 29 HPLC-MS

