

Supplementary Information

Direct acylation of *N*-benzyltriflamides from the alcohol oxidation level via palladium-catalyzed C–H bond activation

Jihye Park,^a Aejin Kim,^a Satyasheel Sharma,^{a,b} Minyoung Kim,^a Eonjeong Park,^a Yukyoung Jeon,^a Youngil Lee,^b Jong Hwan Kwak,^a Young Hoon Jung^a and In Su Kim^{a,*}

^a School of Pharmacy, Sungkyunkwan University, Suwon 440-746, Republic of Korea


^b Department of Chemistry, University of Ulsan, Ulsan 680-749, Republic of Korea.

* Corresponding author. Tel.: +82-31-290-7788; fax: +82-31-292-8800; e-mail: insukim@skku.edu


List of the Contents



¹H NMR and ¹³C NMR copies of all products (3a-k, 4b-l) -----S2–S23

SpinWorks 3: JH 656




SpinWorks 3: JH 656







SpinWorks 3: JH 635




SpinWorks 3: JH 635







SpinWorks 3: JH 632





SpinWorks 3: JH 632















SpinWorks 3: JH 659




SpinWorks 3: JH 659









SpinWorks 3: JH 666




SpinWorks 3: JH 666




SpinWorks 3: JH 667




SpinWorks 3: JH 667




SpinWorks 3: JH 668




SpinWorks 3: JH 668





SpinWorks 3: JH 669




SpinWorks 3: JH 669










SpinWorks 3: JH 673







SpinWorks 3: JH 673














SpinWorks 3: AJ 423



SpinWorks 3: AJ 423



SpinWorks 3: AJ 425



SpinWorks 3: AJ 425

