Supplementary Material

Exploring NMR methods as a tool to select suitable fluorescent nucleotide analogues


Patrick Groves, $*^{a,b}$ Agnieszka Strzelecka-Kiliszek, ^c Anna Sekrecka-Belniak, ^{c,d} Angeles Canales, ^{b,e} Jesús Jiménez-Barbero, ^b Joanna Bandorowicz-Pikula, ^c Sławomir Pikula, ^c and F. Javier Cañada ^b

Supplementary Material

Figure S1. Tr-NOE buildup curves for TNP-GTP binding to AnxA6

Figure S2. Tr-NOE buildup curves for MANT-GTP binding to AnxA6

Figure S3. Tr-NOE buildup curves for BODIPY-γS-GTP binding to AnxA6







Fig. S1. Tr-NOE buildup curves for TNP-GTP binding to AnxA6. The volume of NOE peaks involving proton H8 of the nucleotide base is plotted as a function of mixing time employed in a series of Tr-NOESY experiments. The H8-H1' NOE is represented as filled circles, the H8-H2' NOE is represented by an open circle, the H8-H3' NOE is represented by filled triangles.







Fig. S2. Tr-NOE buildup curves for MANT-GTP binding to AnxA6. The volume of NOE peaks involving proton H8 of the nucleotide base is plotted as a function of mixing time employed in a series of Tr-NOESY experiments. The H8-H1' NOE is represented as filled circles, the H8-H2' NOE is represented by an open circle, the H8-H3' NOE is represented by filled triangles for the O2-MANT isoform. The H8-H1' NOE is represented as open triangles, the H8-H2' NOE is represented by filled squares, the H8-H3' NOE is represented by open squares for the O3-MANT isoform






Fig. S3. Tr-NOE buildup curves for BODIPY-γS-GTP binding to AnxA6. The volume of NOE peaks involving proton H8 of the nucleotide base is plotted as a function of mixing time employed in a series of Tr-NOESY experiments. The H8-H1' NOE is represented as filled circles, the H8-H2' NOE is represented by an open circle, the H8-H3' NOEs were not observed.